

Math 1553 Worksheet §2.8, 2.9

1. Find bases for the column space and the null space of

$$A = \begin{pmatrix} 0 & 1 & -3 & 1 & 0 \\ 1 & -1 & 8 & -7 & 1 \\ -1 & -2 & 1 & 4 & -1 \end{pmatrix}.$$

Solution.

We can row reduce to find that the RREF of $(A \mid 0)$ is

$$\left(\begin{array}{ccccc|c} 1 & 0 & 5 & -6 & 1 & 0 \\ 0 & 1 & -3 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right),$$

so x_3, x_4, x_5 are free, and

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} -5x_3 + 6x_4 - x_5 \\ 3x_3 - x_4 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = x_3 \begin{pmatrix} -5 \\ 3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 6 \\ -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + x_5 \begin{pmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

Therefore, a basis for $\text{Nul } A$ is $\left\{ \begin{pmatrix} -5 \\ 3 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 6 \\ -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \right\}$.

To find a basis for $\text{Col } A$, we use the pivot columns as they were written in the *original* matrix A , *not its RREF*. These are the first two columns:

$$\left\{ \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -2 \end{pmatrix} \right\}.$$

2. Consider the following vectors in \mathbf{R}^3 :

$$b_1 = \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix} \quad b_2 = \begin{pmatrix} 1 \\ 4 \\ 3 \end{pmatrix} \quad u = \begin{pmatrix} 1 \\ 10 \\ 7 \end{pmatrix}$$

Let $V = \text{Span}\{b_1, b_2\}$.

- Explain why $\mathcal{B} = \{b_1, b_2\}$ is a basis for V .
- Determine if u is in V .
- Find a vector b_3 such that $\{b_1, b_2, b_3\}$ is a basis of \mathbf{R}^3 .

Solution.

- a) A quick check shows that b_1 and b_2 are linearly independent (verify!), and we already know they span V , so $\{b_1, b_2\}$ is a basis for V .
- b) u is in V if and only if $c_1 b_1 + c_2 b_2 = u$ for some c_1 and c_2 (in which case $[u]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$ looking ahead to problem 5(b)). We form the augmented matrix $(b_1 \ b_2 \mid u)$ and see if the system is consistent.

$$\left(\begin{array}{cc|c} 2 & 1 & 1 \\ 2 & 4 & 10 \\ 2 & 3 & 7 \end{array} \right) \xrightarrow[\substack{R_2=R_2-R_1 \\ R_3=R_3-R_1}]{R_2=R_2-R_1} \left(\begin{array}{cc|c} 2 & 1 & 1 \\ 0 & 3 & 9 \\ 0 & 2 & 6 \end{array} \right) \xrightarrow[\substack{R_2=R_2/3 \\ R_3=R_3-\frac{2}{3}R_2}]{R_3=R_3-\frac{2}{3}R_2} \left(\begin{array}{cc|c} 2 & 1 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right)$$

The right column is not a pivot column, so the system is consistent, therefore u is in $\text{Span}\{b_1, b_2\}$.

- c) By the increasing span criterion, if we choose b_3 which is not in $\text{Span}\{b_1, b_2\}$, then $\text{Span}\{b_1, b_2, b_3\}$ will be strictly larger than the 2-plane V , so it will be a 3-plane within \mathbf{R}^3 . In other words, the span will be all of \mathbf{R}^3 .

We could choose $b_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, since $(b_1 \ b_2 \ b_3)$ is inconsistent below.

$$\left(\begin{array}{cc|c} 2 & 1 & 1 \\ 2 & 4 & 0 \\ 2 & 3 & 0 \end{array} \right) \xrightarrow[\substack{R_2=R_2-R_1 \\ R_3=R_3-R_1}]{R_2=R_2-R_1} \left(\begin{array}{cc|c} 2 & 1 & 0 \\ 0 & 3 & -1 \\ 0 & 2 & -1 \end{array} \right) \xrightarrow{R_3=R_3-\frac{2}{3}R_2} \left(\begin{array}{cc|c} 2 & 1 & 1 \\ 0 & 3 & -1 \\ 0 & 0 & -1/3 \end{array} \right).$$

3. For (a) and (b), answer “yes” if the statement is always true, “no” if it is always false, and “maybe” otherwise.
- a) If A is an $n \times n$ matrix and $\text{Col } A = \mathbf{R}^n$, then $Ax = 0$ has a nontrivial solution.
- b) If A is an $m \times n$ matrix and $Ax = 0$ has only the trivial solution, then the columns of A form a basis for \mathbf{R}^m .
- c) Give an example of 2×2 matrix whose column space is the same as its null space.

Solution.

- a) No. Since $\text{Col}(A) = \mathbf{R}^n$, the linear transformation $T(x) = Ax$ from \mathbf{R}^n to \mathbf{R}^n is onto, hence T is one-to-one, so $Ax = 0$ has only the trivial solution.
- b) Maybe. If $Ax = 0$ has only the trivial solution and $m = n$, then A is invertible, so the columns of A are linearly independent and span \mathbf{R}^m .

If $m > n$ then the statement is false. For example, $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$ has only the trivial solution for $Ax = 0$, but its columns form only a 2-plane within \mathbf{R}^3 .

- c) Take $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Its null space and column space are $\text{Span}\left\{\begin{pmatrix} 1 \\ 0 \end{pmatrix}\right\}$.

4. In each case, determine whether the given set is a subspace of \mathbf{R}^4 . If it is a subspace, justify why. If it is not a subspace, state a subspace property that it fails.

$$\text{a) } V = \left\{ \begin{pmatrix} x \\ y \\ z \\ w \end{pmatrix} \text{ in } \mathbf{R}^4 \mid x + y = 0 \text{ and } z + w = 0 \right\}$$

$$\text{b) } W = \left\{ \begin{pmatrix} x \\ y \\ z \\ w \end{pmatrix} \text{ in } \mathbf{R}^4 \mid xy - zw = 0 \right\}$$

Solution.

- a) The null space of a 2×4 matrix is automatically a subspace of \mathbf{R}^4 , and V is equal to the null space of the matrix below, so V is a subspace of \mathbf{R}^4 :

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Alternatively, we can verify the subspace properties:

- (1) The zero vector is in V , since $0 + 0 = 0$ and $0 + 0 = 0$.

$$(2) \text{ If } u = \begin{pmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{pmatrix} \text{ and } v = \begin{pmatrix} x_2 \\ y_2 \\ z_2 \\ w_2 \end{pmatrix} \text{ are in } V. \text{ Compute } u + v = \begin{pmatrix} x_1 + x_2 \\ y_1 + y_2 \\ z_1 + z_2 \\ w_1 + w_2 \end{pmatrix}.$$

Are $(x_1 + x_2) + (y_1 + y_2) = 0$ and $(z_1 + z_2) + (w_1 + w_2) = 0$? Yes.

$$(x_1 + x_2) + (y_1 + y_2) = (x_1 + y_1) + (x_2 + y_2) = 0 + 0 = 0,$$

$$(z_1 + z_2) + (w_1 + w_2) = (z_1 + w_1) + (z_2 + w_2) = 0 + 0 = 0.$$

- (3) If $u = \begin{pmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{pmatrix}$ is in V then so is cu for any scalar:

$$cx_1 + cy_1 = c(x_1 + y_1) = c(0) = 0, \quad cz_1 + cw_1 = c(z_1 + w_1) = c(0) = 0.$$

An alternative justification:

- b) Not a subspace. Note $u = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$ and $v = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$ are in W , but $u + v$ is not in W .

$$u + v = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \quad 1 \cdot 1 - 0 \cdot 0 = 1 \neq 0. \quad (W \text{ is not closed under addition})$$

5. This problem covers section 2.9. Parts (a), (b), and (c) are unrelated to each other.

a) True or false: If A is a 3×100 matrix of rank 2, then $\dim(\text{Nul}A) = 97$.

b) For u and \mathcal{B} from problem 2, find $[u]_{\mathcal{B}}$ (the \mathcal{B} -coordinates of u).

c) Let $\mathcal{D} = \left\{ \begin{pmatrix} -2 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$, and suppose $[x]_{\mathcal{D}} = \begin{pmatrix} -1 \\ 3 \end{pmatrix}$. Find x .

Solution.

a) No. By the Rank Theorem, $\text{rank}(A) + \dim(\text{Nul}A) = 100$, so $\dim(\text{Nul}A) = 98$.

b) u is in V if and only if $c_1 b_1 + c_2 b_2 = u$ for some c_1 and c_2 , in which case

$[u]_{\mathcal{B}} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}$. We form the augmented matrix $(b_1 \ b_2 \mid u)$ and solve:

$$\left(\begin{array}{cc|c} 2 & 1 & 1 \\ 2 & 4 & 10 \\ 2 & 3 & 7 \end{array} \right) \xrightarrow[\begin{array}{l} R_2=R_2-R_1 \\ R_3=R_3-R_1 \end{array}]{R_2=R_2-R_1} \left(\begin{array}{cc|c} 2 & 1 & 1 \\ 0 & 3 & 9 \\ 0 & 2 & 6 \end{array} \right) \xrightarrow[\begin{array}{l} R_2=R_2/3 \\ R_3=R_3-\frac{2}{3}R_2 \end{array}]{R_3=R_3-\frac{2}{3}R_2} \left(\begin{array}{cc|c} 2 & 1 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right) \xrightarrow[\text{then } R_1=R_1/2]{R_1=R_1-R_2} \left(\begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{array} \right).$$

We found $c_1 = -1$ and $c_2 = 3$. This means $-b_1 + 3b_2 = u$, so $[u]_{\mathcal{B}} = \begin{pmatrix} -1 \\ 3 \end{pmatrix}$.

c) From $[x]_{\mathcal{D}} = \begin{pmatrix} -1 \\ 3 \end{pmatrix}$, we have

$$x = -d_1 + 3d_2 = -\begin{pmatrix} -2 \\ 1 \end{pmatrix} + 3\begin{pmatrix} 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix} + \begin{pmatrix} 9 \\ 3 \end{pmatrix} = \begin{pmatrix} 11 \\ 2 \end{pmatrix}.$$