

Announcements: Oct 25

- Come pick up your midterm up front
- No WebWork due this week
- No Quiz Friday
- Midterm 3 on Friday Nov 17
- Upcoming Office Hours
 - ▶ Me: Wednesday 3-4, Skiles 234
 - ▶ Qianli: Wednesday 1-2, Clough 280
 - ▶ Arjun: Wednesday, 2:30-3:30, Skiles 230
 - ▶ Kemi: Thursday 9:30-10:30, Skiles 230
 - ▶ Martin: Friday 2-3, Skiles 230

Other sources of help:

- ▶ Math Lab, Clough 280, Mon - Thu 12-6
- ▶ Tutoring: <http://www.successprograms.gatech.edu/tutoring>

(Section 3.1)

Determinants

Outline

- A recursive formula for the determinant
- Special formulas for 2×2 and 3×3
- Inverses via determinants

A formula for the determinant

We will give a **recursive** formula.

First some terminology:

A_{ij} = ij th **minor** of A
= $(n-1) \times (n-1)$ matrix obtained by deleting the i th row and j th column

C_{ij} = $(-1)^{i+j} \det(A_{ij})$
= ij th **cofactor** of A

Finally:

$$\det(A) = a_{11}C_{11} + a_{12}C_{12} + \cdots + a_{1n}C_{1n}$$

A formula for the determinant

The recursive formula:

$$\det(A) = a_{11}C_{11} + a_{12}C_{12} + \cdots + a_{1n}C_{1n}$$

Need to start somewhere...

1×1 matrices

$$\det(a_{11}) =$$

2×2 matrices

$$\det \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} =$$

A formula for the determinant

3×3 matrices

$$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} =$$

Determinants

Compute

$$\det \begin{pmatrix} 5 & 1 & 0 \\ -1 & 3 & 2 \\ 4 & 0 & -1 \end{pmatrix}$$

A formula for the determinant

Another formula for 3×3 matrices

$$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ - a_{13}a_{22}a_{31} - a_{12}a_{21}a_{33} - a_{11}a_{23}a_{32}$$

Use this formula to compute

$$\det \begin{pmatrix} 5 & 1 & 0 \\ -1 & 3 & 2 \\ 4 & 0 & -1 \end{pmatrix}$$

Expanding across other rows and columns

The formula we gave for $\det(A)$ is the **expansion across the first row**. It turns out you can compute the determinant by expanding across any row or column:

$$\det(A) = a_{i1}C_{i1} + \cdots + a_{in}C_{in} \text{ for any fixed } i$$

$$\det(A) = a_{1j}C_{1j} + \cdots + a_{nj}C_{nj} \text{ for any fixed } j$$

Compute:

$$\det \begin{pmatrix} 2 & 1 & 0 \\ 1 & 1 & 0 \\ 5 & 9 & 1 \end{pmatrix}$$

Determinants of triangular matrices

If A is upper (or lower) triangular, $\det(A)$ is easy to compute:

$$\det \begin{pmatrix} 2 & 1 & 5 & -2 \\ 0 & 1 & 2 & -3 \\ 0 & 0 & 5 & 9 \\ 0 & 0 & 0 & 10 \end{pmatrix}$$

Determinants

Poll

What is the determinant?

$$\det \begin{pmatrix} 0 & 7 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \end{pmatrix}$$

A formula for the inverse

(from Section 3.3)

2×2 matrices

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \rightsquigarrow A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

$n \times n$ matrices

$$\begin{aligned} A^{-1} &= \frac{1}{\det(A)} \begin{pmatrix} C_{11} & \cdots & C_{n1} \\ \vdots & \ddots & \vdots \\ C_{1n} & \cdots & C_{nn} \end{pmatrix} \\ &= \frac{1}{\det(A)} (C_{ij})^T \end{aligned}$$

Check that these agree!

The proof uses Cramer's rule (see the notes on the course home page).

A formula for the inverse

(from Section 3.3)

$n \times n$ matrices

$$\begin{aligned} A^{-1} &= \frac{1}{\det(A)} \begin{pmatrix} C_{11} & \cdots & C_{n1} \\ \vdots & \ddots & \vdots \\ C_{1n} & \cdots & C_{nn} \end{pmatrix} \\ &= \frac{1}{\det(A)} (C_{ij})^T \end{aligned}$$

Compute:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}^{-1}$$

Summary

- Special formulas for 2×2 and 3×3 matrices
- Cofactor expansion (works great when there is a row with lots of zeros)
- Formula for the inverse