

Announcements: October 1

- **Midsemester Course Evaluations due Friday** on Canvas
- Midterm 2 Oct 19 in recitation
- **Quiz** on 3.7, 3.9, 4.1 Friday in recitation
- **WeBWork** due Wednesday
- My office hours **Wed 2-3** and Friday 9:30-10:30 in Skiles 234
- TA Office Hours
 - ▶ Arjun Wed 3-4 Skiles 230
 - ▶ Talha Tue/Thu 11-12 Clough 248
 - ▶ Athreya Tue 3-4 Skiles 230
 - ▶ Olivia Thu 3-4 Skiles 230
 - ▶ James Fri 12-1 Skiles 230
 - ▶ Jesse Wed 9:30-10:30 Skiles 230
 - ▶ Vajraang Thu 9:30-10:30 Skiles 230
 - ▶ Hamed Thu 11:15-12, 1:45-2:45, 3-4:15 Clough 280
- Math Lab Monday-Thursday 11:15-5:15 Clough 280
- PLUS Sessions
 - ▶ Tue/Thu 6-7 Clough 280
 - ▶ Mon/Wed 7-8 Clough 123
- Supplemental problems and practice exams on master course web site

Sections 4.2

One-to-one and onto transformations

Section 4.2 Outline

- Learn the definitions of one-to-one and onto functions
- Determine if a given matrix transformation is one-to-one and/or onto

One-to-one

$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** if each b in \mathbb{R}^m is the output for at most one v in \mathbb{R}^n .

In other words: different inputs have different outputs.

Theorem. Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation with matrix A . Then the following are all equivalent:

- T is one-to-one
- the columns of A are linearly independent
- $Ax = 0$ has only the trivial solution
- A has a pivot in each column
- the range of T has dimension n

What can we say about the relative sizes of m and n if T is one-to-one?

Draw a picture of the range of a one-to-one mapping $\mathbb{R} \rightarrow \mathbb{R}^3$.

Onto

$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** if the range of T equals the codomain \mathbb{R}^m , that is, each b in \mathbb{R}^m is the output for at least one input v in \mathbb{R}^n .

Theorem. Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation with matrix A . Then the following are all equivalent:

- T is onto
- the columns of A span \mathbb{R}^m
- A has a pivot in each row
- $Ax = b$ is consistent for all b in \mathbb{R}^m
- the range of T has dimension m

What can we say about the relative sizes of m and n if T is onto?

Give an example of an onto mapping $\mathbb{R}^3 \rightarrow \mathbb{R}$.

One-to-one and Onto

Do the following give matrix transformations that are one-to-one? onto?

$$\begin{pmatrix} 1 & 0 & 7 \\ 0 & 1 & 2 \\ 0 & 0 & 9 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$$

One-to-one and Onto

Which of the previously-studied matrix transformations of \mathbb{R}^2 are one-to-one? Onto?

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ reflection}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \text{ projection}$$

$$\begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix} \text{ scaling}$$

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ shear}$$

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \text{ rotation}$$

Summary of Section 4.2

- $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **one-to-one** if each b in \mathbb{R}^m is the output for at most one v in \mathbb{R}^n .
- **Theorem.** Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation with matrix A . Then the following are all equivalent:
 - ▶ T is one-to-one
 - ▶ the columns of A are linearly independent
 - ▶ $Ax = 0$ has only the trivial solution
 - ▶ A has a pivot in each column
 - ▶ the range has dimension n
- $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **onto** if the range of T equals the codomain \mathbb{R}^m , that is, each b in \mathbb{R}^m is the output for at least one input v in \mathbb{R}^n .
- **Theorem.** Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation with matrix A . Then the following are all equivalent:
 - ▶ T is onto
 - ▶ the columns of A span \mathbb{R}^m
 - ▶ A has a pivot in each row
 - ▶ $Ax = b$ is consistent for all b in \mathbb{R}^m .
 - ▶ the range of T has dimension m