

## Announcements: October 24

- Come pick up your midterm up front.
- Midterm 3 on §4.5-6.5 **Nov 16** in recitation
- **No Quiz** Friday in recitation
- **WeBWork** 4.5 due **Wednesday**
- My office hours **Wed 2-3** and Friday 9:30-10:30 in Skiles 234
- TA office hours
  - ▶ Arjun Wed 3-4 Skiles 230
  - ▶ Talha Tue/Thu 11-12 Clough 248
  - ▶ Athreya Tue 3-4 Skiles 230
  - ▶ Olivia Thu 3-4 Skiles 230
  - ▶ James Fri 12-1 Skiles 230
  - ▶ Jesse Wed 9:30-10:30 Skiles 230
  - ▶ Vajraang Thu 9:30-10:30 Skiles 230
  - ▶ Hamed Thu 11:15-12, 1:45-2:45, 3-4:15 Clough 280
- Math Lab Monday-Thursday 11:15-5:15 Clough 280 [▶ Schedule](#)
- PLUS Sessions
  - ▶ Tue/Thu 6-7 Clough 280
  - ▶ Mon/Wed 7-8 Clough 123

## Where are we?

- We have studied the problem  $Ax = b$
- We next want to study  $Ax = \lambda x$
- At the end of the course we want to almost solve  $Ax = b$

We need determinants for the second item.

# Sections 5.1 and 5.3

The definition of the determinant and volumes


## Outline of Sections 5.1 and 5.3

- Volume and invertibility
- A definition of determinant in terms of row operations
- Using the definition of determinant to compute the determinant
- Determinants of products:  $\det(AB)$
- Determinants and linear transformations and volumes

## Invertibility and volume


When is a  $2 \times 2$  matrix invertible?

When the rows (or columns) don't lie on a line  $\Leftrightarrow$  the corresponding parallelogram has non-zero area



When is a  $3 \times 3$  matrix invertible?

When the rows (or columns) don't lie on a plane  $\Leftrightarrow$  the corresponding parallelepiped (3D parallelogram) has non-zero volume



Same for  $n \times n$ !

## The definition of determinant

The **determinant** of a *square* matrix is a number so that

1. If we do a row replacement on a matrix, the determinant is unchanged
2. If we swap two rows of a matrix, the determinant scales by  $-1$
3. If we scale a row of a matrix by  $k$ , the determinant scales by  $k$
4.  $\det(I_n) = 1$

Why would we think of this? *Answer: This is exactly how volume works.*

Try it out for  $2 \times 2$  matrices.

## The definition of determinant

The **determinant** of a *square* matrix is a number so that

1. If we do a row replacement on a matrix, the determinant is unchanged
2. If we swap two rows of a matrix, the determinant scales by  $-1$
3. If we scale a row of a matrix by  $k$ , the determinant scales by  $k$
4.  $\det(I_n) = 1$

*Problem.* Just using these rules, compute the determinants:

$$\begin{pmatrix} 1 & 0 & 8 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & 17 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{pmatrix}$$

## A basic fact about determinants

**Fact.** If  $A$  has a zero row, then  $\det(A) = 0$ .

Why does this follow from the definition?



## A first formula for the determinant

**Fact.** Suppose we row reduce  $A$ . Then

$$\det A = (-1)^{\#\text{row swaps used}} \left( \frac{\text{product of diagonal entries of row reduced matrix}}{\text{product of scalings used}} \right)$$

What is the determinant of a matrix in row echelon form?

Use the fact to get a formula for the determinant of any  $2 \times 2$  matrix.

Consequence of the above fact:

**Fact.**  $\det A \neq 0 \Leftrightarrow A$  invertible

## Computing determinants

...using the definition in terms of row operations

$$\det \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 5 & 7 & -4 \end{pmatrix} =$$

## A Mathematical Conundrum

We have this definition of a determinant, and it gives us a way to compute it.

But: we don't know that such a determinant function exists.

More specifically, we haven't ruled out the possibility that two different row reductions might give us two different answers for the determinant.

Don't worry! It is all okay.

We already gave the key idea: that determinant is just the volume of the corresponding parallelepiped. You can read the proof in the book if you want.

**Fact 1.** There is such a number  $\det$  and it is unique.

## Properties of the determinant

Fact 1. There is such a number  $\det$  and it is unique.

Fact 2.  $A$  is invertible  $\Leftrightarrow \det(A) \neq 0$       **important!**

Fact 3.  $\det A = (-1)^{\#\text{row swaps used}} \left( \frac{\text{product of diagonal entries of row reduced matrix}}{\text{product of scalings used}} \right)$

Fact 4. The function can be computed by any of the  $2n$  cofactor expansions.

Fact 5.  $\det(AB) = \det(A) \det(B)$       **important!**

Fact 6.  $\det(A^T) = \det(A)$       **ok, now we need to say what transpose is**

Fact 7.  $\det(A)$  is signed volume of the parallelepiped spanned by cols of  $A$ .

If you want the proofs, see the book. Actually Fact 1 is the hardest!

## Powers

**Fact 5.**  $\det(AB) = \det(A) \det(B)$

Use this fact to compute

$$\det \left( \left( \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 5 & 7 & -4 \end{pmatrix} \right)^5 \right)$$

What is  $\det(A^{-1})$ ?

## Poll

Suppose we know  $A^5$  is invertible. Is  $A$  invertible?

1. yes
2. no
3. maybe

## Areas of triangles

What is the area of the triangle in  $\mathbb{R}^2$  with vertices  $(1, 2)$ ,  $(4, 3)$ , and  $(2, 5)$ ?


What is the area of the parallelogram in  $\mathbb{R}^2$  with vertices  $(1, 2)$ ,  $(4, 3)$ ,  $(2, 5)$ , and  $(5, 6)$ ?

## Determinants and linear transformations

Say  $A$  is an  $n \times n$  matrix and  $T(v) = Av$ .

**Fact 8.** If  $S$  is some subset of  $\mathbb{R}^n$ , then  $\text{vol}(T(S)) = |\det(A)| \cdot \text{vol}(S)$ .

This works even if  $S$  is curvy, like a circle or an ellipse, or:



Why? First check it for little squares/cubes (Fact 7). Then: Calculus!



## Summary of Sections 5.1 and 5.3

Say  $\det$  is a function  $\det : \{\text{matrices}\} \rightarrow \mathbb{R}$  with:

1.  $\det(I_n) = 1$
2. If we do a row replacement on a matrix, the determinant is unchanged
3. If we swap two rows of a matrix, the determinant scales by  $-1$
4. If we scale a row of a matrix by  $k$ , the determinant scales by  $k$

**Fact 1.** There is such a function  $\det$  and it is unique.

**Fact 2.**  $A$  is invertible  $\Leftrightarrow \det(A) \neq 0$       **important!**

**Fact 3.**  $\det A = (-1)^{\#\text{row swaps used}} \left( \frac{\text{product of diagonal entries of row reduced matrix}}{\text{product of scalings used}} \right)$

**Fact 4.** The function can be computed by any of the  $2n$  cofactor expansions.

**Fact 5.**  $\det(AB) = \det(A) \det(B)$       **important!**

**Fact 6.**  $\det(A^T) = \det(A)$

**Fact 7.**  $\det(A)$  is signed volume of the parallelepiped spanned by cols of  $A$ .

**Fact 8.** If  $S$  is some subset of  $\mathbb{R}^n$ , then  $\text{vol}(T(S)) = |\det(A)| \cdot \text{vol}(S)$ .