

## Announcements: November 5

- Midterm 3 on §4.5-6.5 **Nov 16** in recitation
- **Quiz** 6.1-6.2 Friday in recitation.
- **WeBWork** 6.1-6.2 due **Wednesday**
- My office hours **Wed 2-3** and Friday 9:30-10:30 in Skiles 234
- TA office hours
  - ▶ Arjun Wed 3-4 Skiles 230
  - ▶ Talha Tue/Thu 11-12 Clough 248
  - ▶ Athreya Tue 3-4 Skiles 230
  - ▶ Olivia Thu 3-4 Skiles 230
  - ▶ James Tue 11-12 Skiles 230
  - ▶ Jesse Wed 9:30-10:30 Skiles 230
  - ▶ Vajraang Thu 11-12 Skiles 230
  - ▶ Hamed Thu 11:15-12, 1:45-2:45, 3-4:15 Clough 280
- Math Lab Monday-Thursday 11:15-5:15 Clough 280 [▶ Schedule](#)
- PLUS Sessions
  - ▶ Tue/Thu 6-7
  - ▶ Mon/Wed 6-7

# Eigenvalues in Structural Engineering

Watch this video about the Tacoma Narrows bridge. [▶ Watch](#)

Here are some toy models. [▶ Check it out](#)

The masses move the most at their **natural frequencies**  $\omega$ . To find those, use the spring equation:  $mx'' = -kx \rightsquigarrow \sin(\omega t)$ .

With 3 springs and 2 equal masses, we get:

$$mx_1'' = -kx_1 + k(x_2 - x_1)$$

$$mx_2'' = -kx_2 + k(x_1 - x_2)$$

Guess a solution  $x_1(t) = A_1(\cos(\omega t) + i \sin(\omega t))$  and similar for  $x_2$ . Finding  $\omega$  reduces to finding **eigenvalues** of  $\begin{pmatrix} -2k & k \\ k & -2k \end{pmatrix}$ .

Eigenvectors:  $(1, 1)$  &  $(1, -1)$  (in/out of phase) [▶ Details](#)

# Section 6.4

## Diagonalization

## Section 6.4 Outline

- Diagonalization
- Using diagonalization to take powers
- Algebraic versus geometric dimension

## We understand diagonal matrices

We completely understand what diagonal matrices do to  $\mathbb{R}^n$ . For example:

$$\begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$$

We have seen that it is useful to take powers of matrices: for instance in computing rabbit populations.

If  $A$  is diagonal, powers of  $A$  are easy to compute. For example:

$$\begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}^{10} = \dots$$

## Powers of matrices that are similar to diagonal ones

What if  $A$  is not diagonal? Suppose want to understand the matrix

$$A = \begin{pmatrix} 5/4 & 3/4 \\ 3/4 & 5/4 \end{pmatrix}$$

geometrically? Or take it's 10th power? What would we do?

What if I give you the following equality:

$$\begin{pmatrix} 5/4 & 3/4 \\ 3/4 & 5/4 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}^{-1}$$

$A \qquad = \qquad C \qquad D \qquad C^{-1}$

This is called **diagonalization**.

How does this help us understand  $A$ ? Or find  $A^{10}$ ? [▶ Demo](#)

# Diagonalization

Suppose  $A$  is  $n \times n$ . We say that  $A$  is **diagonalizable** if we can write:

$$A = CDC^{-1} \quad D = \text{diagonal}$$

We say that  $A$  is similar to  $D$ .

How does this factorization of  $A$  help describe what  $A$  **does** to  $\mathbb{R}^n$ ?  
How does this help us take powers of  $A$ ?

Understanding the rabbit example: since 2 is the largest eigenvalue, (almost) all other vectors get pulled towards that eigenvector. Compare with the example from the last slide.

# Diagonalization

**Theorem.**  $A$  is diagonalizable  $\Leftrightarrow A$  has  $n$  linearly independent eigenvectors.

In this case

$$A = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix} \begin{pmatrix} \lambda_1 & & & \\ & \ddots & & \\ & & \lambda_n & \\ & & & \end{pmatrix} \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix}^{-1}$$
$$= \quad \quad \quad C \quad \quad \quad D \quad \quad \quad C^{-1}$$

where  $v_1, \dots, v_n$  are linearly independent eigenvectors and  $\lambda_1, \dots, \lambda_n$  are the corresponding eigenvalues (in **order**).

Why?



## Example

Diagonalize if possible.

$$\begin{pmatrix} 2 & 6 \\ 0 & -1 \end{pmatrix}$$

## Example

Diagonalize if possible.

$$\begin{pmatrix} 3 & 1 \\ 0 & 3 \end{pmatrix}$$

## Example

Diagonalize if possible.

$$\begin{pmatrix} 3/4 & 1/4 \\ 1/4 & 3/4 \end{pmatrix}$$

▶ Demo

## More Examples

Diagonalize if possible.

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} a & b \\ b & a \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix}$$

# Poll

Poll

Which are diagonalizable?

$$\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix} \quad \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

## Distinct Eigenvalues

Fact. If  $A$  has  $n$  distinct eigenvalues, then  $A$  is diagonalizable.

Why?

# Non-Distinct Eigenvalues

Theorem. Suppose

- $A = n \times n$ , has eigenvalues  $\lambda_1, \dots, \lambda_k$
- $a_i =$  algebraic multiplicity of  $\lambda_i$
- $d_i =$  dimension of  $\lambda_i$  eigenspace (“geometric multiplicity”)

Then

1.  $d_i \leq a_i$  for all  $i$
2.  $A$  is diagonalizable  $\Leftrightarrow \sum d_i = n$   
 $\Leftrightarrow \sum a_i = n$  and  $d_i = a_i$  for all  $i$

So: if you find one eigenvalue where the geometric multiplicity is less than the algebraic multiplicity, the matrix is not diagonalizable.

## Summary of Section 6.4

- $A$  is diagonalizable if  $A = CDC^{-1}$  where  $D$  is diagonal
- A diagonal matrix stretches along its eigenvectors by the eigenvalues, similar to a diagonal matrix
- If  $A = CDC^{-1}$  then  $A^k = CD^kC^{-1}$
- $A$  is diagonalizable  $\Leftrightarrow A$  has  $n$  linearly independent eigenvectors  $\Leftrightarrow$  the sum of the geometric dimensions of the eigenspaces in  $n$
- If  $A$  has  $n$  distinct eigenvalues it is diagonalizable