

Math 1553 Worksheet §5.4, 5.5

1. Answer yes, no, or maybe. Justify your answers. In each case, A is a matrix whose entries are real numbers.
- a) If A is a 3×3 matrix with characteristic polynomial $-\lambda(\lambda - 5)^2$, then the 5-eigenspace is 2-dimensional.
 - b) If A is an invertible 2×2 matrix, then A is diagonalizable.
 - c) A 3×3 matrix A can have a non-real complex eigenvalue with multiplicity 2.

Solution.

- a) Maybe. The geometric multiplicity of $\lambda = 5$ can be 1 or 2. For example, the matrix $\begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ has a 5-eigenspace which is 2-dimensional, whereas the matrix $\begin{pmatrix} 5 & 1 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ has a 5-eigenspace which is 1-dimensional. Both matrices have characteristic polynomial $-\lambda(5 - \lambda)^2$.
- b) Maybe. The identity matrix is invertible and diagonalizable, but the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ is invertible but not diagonalizable.
- c) No. If c is a (non-real) complex eigenvalue with multiplicity 2, then its conjugate \bar{c} is an eigenvalue with multiplicity 2 since complex eigenvalues always occur in conjugate pairs. This would mean A has a characteristic polynomial of degree 4 or more, which is impossible for a 3×3 matrix.

2. $A = \begin{pmatrix} 2 & 3 & 1 \\ 3 & 2 & 4 \\ 0 & 0 & -1 \end{pmatrix}.$

a) Find the eigenvalues of A , and find a basis for each eigenspace.

b) Is A diagonalizable? If your answer is yes, find a diagonal matrix D and an invertible matrix C so that $A = CDC^{-1}$. If your answer is no, justify why A is not diagonalizable.

Solution.

a) We solve $0 = \det(A - \lambda I)$.

$$0 = \det \begin{pmatrix} 2-\lambda & 3 & 1 \\ 3 & 2-\lambda & 4 \\ 0 & 0 & -1-\lambda \end{pmatrix} = (-1-\lambda)(-1)^6 \det \begin{pmatrix} 2-\lambda & 3 \\ 3 & 2-\lambda \end{pmatrix} = (-1-\lambda)((2-\lambda)^2 - 9)$$

$$= (-1-\lambda)(\lambda^2 - 4\lambda - 5) = -(\lambda + 1)^2(\lambda - 5).$$

So $\lambda = -1$ and $\lambda = 5$ are the eigenvalues.

$$\underline{\lambda = -1}: (A + I | 0) = \left(\begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 3 & 3 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow{R_2=R_2-R_1} \left(\begin{array}{ccc|c} 3 & 3 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow[\text{then } R_1=R_1/3]{R_1=R_1-R_2}$$

$$\left(\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right), \text{ with solution } x_1 = -x_2, x_2 = x_2, x_3 = 0. \text{ The } (-1)\text{-eigenspace}$$

has basis $\left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \right\}.$

$\lambda = 5$:

$$(A - 5I | 0) = \left(\begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 3 & -3 & 4 & 0 \\ 0 & 0 & -6 & 0 \end{array} \right) \xrightarrow[\text{then } R_2 \leftrightarrow R_3, R_1/(-3)]{\substack{R_2=R_2+R_1 \\ R_3=R_3/(-6)}} \left(\begin{array}{ccc|c} -3 & 3 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right) \xrightarrow{R_1=R_1-R_3, R_2=R_2-5R_3} \left(\begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$$

with solution $x_1 = x_2, x_2 = x_2, x_3 = 0$. The 5-eigenspace has basis $\left\{ \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}.$

b) A is a 3×3 matrix that only admits 2 linearly independent eigenvectors, so A is not diagonalizable.