

IMPORTANT DEFINITIONS AND THEOREMS
REFERENCE SHEET

This is a (not quite comprehensive) list of definitions and theorems given in Math 1553. Pay particular attention to the ones in **red**.

Study Tip

For each definition, find an example of something that satisfies the requirements of the definition, and an example of something that does not. For each theorem, find an example of something that satisfies the hypotheses of the theorem, and an example of something that does not satisfy the conclusions (or the hypotheses, of course) of the theorem. This is *great* conceptual practice.

CHAPTER 2

SECTION 2.1.

Definition. \mathbf{R}^n = all ordered n -tuples of real numbers $(x_1, x_2, x_3, \dots, x_n)$.

This is the number line when $n = 1$, the xy -plane when $n = 2$, and space when $n = 3$.

Definition. A **solution** to a system of linear equations is a list of numbers making *all* of the equations true.

A solution of a system of equations in n variables is a point in \mathbf{R}^n .

Definition. The **elementary row operations** are the following matrix operations:

- Multiply all entries in a row by a nonzero number (scale).
- Add (a multiple of) each entry of one row to the corresponding entry in another (row replacement).
- Swap two rows.

Definition. Two matrices are called **row equivalent** if one can be obtained from the other by doing some number of elementary row operations.

Definition. A system of equations is called **inconsistent** if it has no solution. It is **consistent** otherwise.

SECTION 2.2.

Definition. A matrix is in **row echelon form** if

- (1) All zero rows are at the bottom.

- (2) Each leading nonzero entry of a row is to the right of the leading entry of the row above.
- (3) Below a leading entry of a row, all entries are zero.

Definition. A **pivot** is the first nonzero entry of a row of a matrix in row echelon form.

Definition. A matrix is in **reduced row echelon form** if it is in row echelon form, and in addition,

- (4) The pivot in each nonzero row is equal to 1.
- (5) Each pivot is the only nonzero entry in its column.

Theorem. *Every matrix is row equivalent to one and only one matrix in reduced row echelon form.*

Review. **Row reduction algorithm.**

SECTION 2.3.

Definition. Consider a *consistent* linear system of equations in the variables x_1, \dots, x_n . Let A be the reduced row echelon form of the matrix for this system. We say that x_i is a **free variable** if its corresponding column in A is *not* a pivot column.

Definition. The **parametric form** for the general solution to a system of equations is a system of equations for the non-free variables in terms of the free variables. For instance, if x_2 and x_4 are free,

$$x_1 = 2 - 3x_4 \quad x_3 = -1 - 4x_4$$

is a parametric form.

Theorem. *Every solution to a consistent linear system is obtained by substituting (unique) values for the free variables in the parametric form.*

Fact. *There are three possibilities for the solution set of a linear system with augmented matrix A :*

- (1) *The system is inconsistent: it has zero solutions, and the last column of A is a pivot column.*
- (2) *The system has a unique solution: every column of A except the last is a pivot column.*
- (3) *The system has infinitely many solutions: the last column isn't a pivot column, and some other column isn't either. These last columns correspond to free variables.*

CHAPTER 3

SECTION 3.1.

Definition. A **vector** is an arrow with a given length and direction.

Definition. A **scalar** is another name for a real number (to distinguish it from a vector).

Review. Parallelogram law for vector addition, geometric interpretation of vector subtraction and scalar multiplication.

Definition. A **linear combination** of vectors v_1, v_2, \dots, v_n is a vector of the form

$$c_1 v_1 + c_2 v_2 + \cdots + c_n v_n$$

where c_1, c_2, \dots, c_n are scalars, called the **weights** or **coefficients** of the linear combination.

SECTION 3.2.

Definition. A **vector equation** is an equation involving vectors. (It is equivalent to a list of equations involving only scalars.)

Definition. The **span** of a set of vectors v_1, v_2, \dots, v_n is the set of all linear combinations of these vectors:

$$\text{Span}\{v_1, \dots, v_p\} = \{x_1 v_1 + \cdots + x_p v_p \mid x_1, \dots, x_p \text{ in } \mathbf{R}\}.$$

Review. Pictures of spans.

Theorem. *The following are equivalent:*

- (1) A vector b is in the span of v_1, v_2, \dots, v_p .
- (2) The vector equation

$$x_1 v_1 + x_2 v_2 + \cdots + x_p v_p$$

has a solution.

- (3) The linear system with augmented matrix

$$\left(\begin{array}{c|c|c|c|c} | & | & \cdots & | & | \\ v_1 & v_2 & & v_p & b \\ | & | & & | & | \end{array} \right)$$

is consistent.

Review. Pictures of consistent and inconsistent vector equations.

SECTION 3.3.

Definition. The **product** of an $m \times n$ matrix A with a vector x in \mathbf{R}^n is the linear combination

$$Ax = \left(\begin{array}{c|c|c|c} | & | & \cdots & | \\ v_1 & v_2 & & v_n \\ | & | & & | \end{array} \right) \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} := x_1 v_1 + x_2 v_2 + \cdots + x_n v_n.$$

The output is a vector in \mathbf{R}^m .

The product can also be computed by multiplying rows:

$$Ax = \begin{pmatrix} -r_1- \\ -r_2- \\ \vdots \\ -r_m- \end{pmatrix} x = \begin{pmatrix} r_1 x \\ r_2 x \\ \vdots \\ r_m x \end{pmatrix}.$$

Definition. A **matrix equation** is a vector equation involving a product of a matrix with a vector.

We now have *four* equivalent ways of writing (and thinking about) linear systems:

(1) As a system of equations

$$\begin{aligned} 2x_1 + 3x_2 &= 7 \\ x_1 - x_2 &= 5 \end{aligned}$$

(2) As an augmented matrix:

$$\left(\begin{array}{cc|c} 2 & 3 & 7 \\ 1 & -1 & 5 \end{array} \right)$$

(3) As a vector equation ($x_1v_1 + \cdots + x_nv_n = b$):

$$x_1 \begin{pmatrix} 2 \\ 1 \end{pmatrix} + x_2 \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

(4) As a matrix equation ($Ax = b$):

$$\begin{pmatrix} 2 & 3 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

In particular, *all four have the same solution set.*

Theorem. $Ax = b$ has a solution if and only if b is in the span of the columns of A .

Theorem. Let A be an $m \times n$ (non-augmented) matrix. The following are equivalent

- (1) $Ax = b$ has a solution for all b in \mathbf{R}^m .
- (2) The span of the columns of A is all of \mathbf{R}^m .
- (3) A has a pivot in each row.

SECTION 3.4.

Definition. A system of linear equations of the form $Ax = 0$ is called **homogeneous**.

Definition. A system of linear equations of the form $Ax = b$ for $b \neq 0$ is called **inhomogeneous** or **non-homogeneous**.

Definition. The **trivial solution** to a homogeneous equation is the solution $x = 0$: $A0 = 0$.

Theorem. Let A be a matrix. The following are equivalent:

- (1) $Ax = 0$ has a nontrivial solution.
- (2) There is a free variable.
- (3) A has a column with no pivot.

Theorem. The solution set of a homogeneous equation $Ax = 0$ is a span. You can find a spanning set by computing the parametric vector form.

Definition. The **parametric vector form** for the general solution to a system of equations $Ax = b$ is a vector equation expressing all variables in terms of the free variables. For instance, if x_2 and x_4 are free,

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \\ -1 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -3 \\ 0 \\ -4 \\ 1 \end{pmatrix}$$

is a parametric vector form. The constant vector $(2, 0, -1, 0)$ is a **specific solution** or **particular solution** to $Ax = b$.

Theorem. *The solution set of a linear system $Ax = b$ is a translate of the solution set of $Ax = 0$ by a specific solution.*

SECTION 3.5.

Definition. A set of vectors $\{v_1, v_2, \dots, v_p\}$ in \mathbf{R}^n is **linearly independent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has only the trivial solution $x_1 = x_2 = \dots = x_p = 0$.

Definition. A set of vectors $\{v_1, v_2, \dots, v_p\}$ in \mathbf{R}^n is **linearly dependent** if the vector equation

$$x_1 v_1 + x_2 v_2 + \dots + x_p v_p = 0$$

has a nontrivial solution (not all x_i are zero). Such a solution is a **linear dependence relation**.

Theorem. *A set of vectors $\{v_1, v_2, \dots, v_p\}$ is linearly dependent if and only if one of the vectors is in the span of the other ones.*

Fact. *Say v_1, v_2, \dots, v_n are in \mathbf{R}^m . If $n > m$ then $\{v_1, v_2, \dots, v_n\}$ is linearly dependent.*

Fact. *If one of v_1, v_2, \dots, v_n is zero, then $\{v_1, v_2, \dots, v_n\}$ is linearly dependent.*

Theorem. *Let v_1, v_2, \dots, v_n be vectors in \mathbf{R}^m , and let A be the $m \times n$ matrix with columns v_1, v_2, \dots, v_n . The following are equivalent:*

- (1) *The set $\{v_1, v_2, \dots, v_n\}$ is linearly independent.*
- (2) *No one vector is in the span of the others.*
- (3) *For every j between 1 and n , v_j is not in $\text{Span}\{v_1, v_2, \dots, v_{j-1}\}$.*
- (4) *$Ax = 0$ only has the trivial solution.*
- (5) *A has a pivot in every column.*

Review. Pictures of linear dependence and independence.

SECTION 3.6.

Definition. A **subspace** of \mathbf{R}^n is a subset V of \mathbf{R}^n satisfying:

- (1) The zero vector is in V .
- (2) If u and v are in V , then $u + v$ is also in V .
- (3) If u is in V and c is in \mathbf{R} , then cu is in V .

Definition. If $V = \text{Span}\{v_1, v_2, \dots, v_n\}$, we say that V is the subspace **generated by** or **spanned by** the vectors v_1, v_2, \dots, v_n .

Theorem. A subspace is a span, and a span is a subspace.

Definition. The **column space** of a matrix A is the subspace spanned by the columns of A . It is written $\text{Col}A$.

Definition. The **null space** of A is the set of all solutions of the homogeneous equation $Ax = 0$:

$$\text{Nul}A = \{x \mid Ax = 0\}.$$

Example. The following are the most important examples of subspaces in this class (some won't appear until later):

- Any $\text{Span}\{v_1, v_2, \dots, v_m\}$.
- The column space of a matrix: $\text{Col}A = \text{Span}\{\text{columns of } A\}$.
- The range of a linear transformation (same as above).
- The null space of a matrix: $\text{Nul}A = \{x \mid Ax = 0\}$.
- The row space of a matrix: $\text{Row}A = \text{Span}\{\text{rows of } A\}$.
- The λ -eigenspace of a matrix, where λ is an eigenvalue.
- The orthogonal complement W^\perp of a subspace W .
- The zero subspace $\{0\}$.
- All of \mathbf{R}^n .

SECTION 3.7.

Definition. Let V be a subspace of \mathbf{R}^n . A **basis** of V is a set of vectors $\{v_1, v_2, \dots, v_m\}$ in V such that:

- (1) $V = \text{Span}\{v_1, v_2, \dots, v_m\}$, and
- (2) $\{v_1, v_2, \dots, v_m\}$ is linearly independent.

The number of vectors in a basis is the **dimension** of V , and is written $\text{dim } V$.

Theorem. Every basis for a subspace has the same number of vectors in it.

Fact. The vectors in the parametric vector form of the general solution to $Ax = 0$ always form a basis for $\text{Nul}A$.

Fact. The pivot columns of A always form a basis for $\text{Col}A$.

SECTION 3.9.

Definition. The **rank** of a matrix A , written $\text{rank}A$, is the dimension of the column space $\text{Col}A$.

Rank Theorem. If A is an $m \times n$ matrix, then

$$\text{rank}A + \dim \text{Nul}A = n = \text{the number of columns of } A.$$

Basis Theorem. Let V be a subspace of dimension m . Then:

- Any m linearly independent vectors in V form a basis for V .
- Any m vectors that span V form a basis for V .

CHAPTER 4

SECTION 4.1.

Definition. A **transformation** (or **function** or **map**) from \mathbf{R}^n to \mathbf{R}^m is a rule T that assigns to each vector x in \mathbf{R}^n a vector $T(x)$ in \mathbf{R}^m .

- \mathbf{R}^n is called the **domain** of T .
- \mathbf{R}^m is called the **codomain** of T .
- For x in \mathbf{R}^n , the vector $T(x)$ in \mathbf{R}^m is the **image** of x under T .
Notation: $x \mapsto T(x)$.
- The set of all images $\{T(x) \mid x \text{ in } \mathbf{R}^n\}$ is the **range** of T .

Notation. $T: \mathbf{R}^n \longrightarrow \mathbf{R}^m$ means T is a transformation from \mathbf{R}^n to \mathbf{R}^m .

Definition. Let A be an $m \times n$ matrix. The **matrix transformation** associated to A is the transformation

$$T: \mathbf{R}^n \longrightarrow \mathbf{R}^m \quad \text{defined by} \quad T(x) = Ax.$$

- The domain is \mathbf{R}^n , where n is the number of columns of A .
- The codomain is \mathbf{R}^m , where m is the number of rows of A .
- The range is the span of the columns of A .

Review. Geometric transformations: **projection, reflection, rotation, dilation, shear.**

SECTION 4.2.

Definition. A transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is **onto** (or **surjective**) if the range of T is equal to \mathbf{R}^m (its codomain). In other words, each b in \mathbf{R}^m is the image of *at least one* x in \mathbf{R}^n .

Theorem. Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- T is onto
- $T(x) = b$ has a solution for every b in \mathbf{R}^m
- $Ax = b$ is consistent for every b in \mathbf{R}^m
- The columns of A span \mathbf{R}^m
- A has a pivot in every row.

Definition. A transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ is **one-to-one** (or **into**, or **injective**) if different vectors in \mathbf{R}^n map to different vectors in \mathbf{R}^m . In other words, each b in \mathbf{R}^m is the image of at most one x in \mathbf{R}^n .

Theorem. Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation with matrix A . Then the following are equivalent:

- T is one-to-one
- $T(x) = b$ has one or zero solutions for every b in \mathbf{R}^m
- $Ax = b$ has a unique solution or is inconsistent for every b in \mathbf{R}^m
- $Ax = 0$ has a unique solution
- The columns of A are linearly independent
- A has a pivot in every column.

SECTION 4.3.

Definition. A **linear transformation** is a transformation T satisfying

$$T(u + v) = T(u) + T(v) \quad \text{and} \quad T(cv) = cT(v)$$

for all vectors u, v and all scalars c .

Definition. The **unit coordinate vectors** in \mathbf{R}^n are

$$e_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \dots, \quad e_{n-1} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad e_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$

Fact. If A is a matrix, then Ae_i is the i th column of A .

Definition. Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear transformation. The **standard matrix** for T is

$$\left(\begin{array}{c|c|c|c} & | & | & \cdots & | \\ T(e_1) & T(e_2) & \cdots & T(e_n) \\ & | & | & & | \end{array} \right).$$

Theorem. If T is a linear transformation, then it is the matrix transformation associated to its standard matrix.

SECTION 4.4.

Definition. The **ij th entry** of a matrix A is the entry in the i th row and j th column. Notation: a_{ij} .

Definition. The entries $a_{11}, a_{22}, a_{33}, \dots$ are the **diagonal entries**; they form the **main diagonal** of the matrix.

Definition. A **diagonal matrix** is a square matrix whose only nonzero entries are on the main diagonal.

Definition. The $n \times n$ **identity matrix** I_n is the diagonal matrix with all diagonal entries equal to 1. It has the property that $I_n A = A$ for any $n \times m$ matrix A .

Definition. The **zero matrix** (of size $m \times n$) is the $m \times n$ matrix 0 with all zero entries.

Definition. The **transpose** of an $m \times n$ matrix A is the $n \times m$ matrix A^T whose rows are the columns of A . In other words, the ij entry of A^T is a_{ji} .

Definition. The **product** of an $m \times n$ matrix A with an $n \times p$ matrix B is the $m \times p$ matrix

$$AB = \left(\begin{array}{c|c|c|c} | & | & & | \\ Av_1 & Av_2 & \cdots & Av_p \\ | & | & & | \end{array} \right),$$

where v_1, v_2, \dots, v_p are the columns of B .

Fact. Suppose A has is an $m \times n$ matrix, and that the other matrices below have the right size to make multiplication work. Then:

$$\begin{array}{ll} A(BC) = (AB)C & A(B + C) = AB + AC \\ (B + C)A = BA + CA & c(AB) = (cA)B \\ c(AB) = A(cB) & I_n A = A \\ AI_m = A & \end{array}$$

Fact. If $A, B,$ and C are matrices, then:

- (1) AB is usually not equal to BA .
- (2) $AB = AC$ does not imply $B = C$.
- (3) $AB = 0$ does not imply $A = 0$ or $B = 0$.

Definition. Let $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U : \mathbf{R}^p \rightarrow \mathbf{R}^n$ be transformations. The **composition** is the transformation

$$T \circ U : \mathbf{R}^p \rightarrow \mathbf{R}^m \quad \text{defined by} \quad T \circ U(x) = T(U(x)).$$

Theorem. Let $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ and $U : \mathbf{R}^p \rightarrow \mathbf{R}^n$ be linear transformations with matrices A and B , respectively. Then the matrix for $T \circ U$ is AB .

SECTION 4.5.

Definition. A square matrix A is **invertible** (or **nonsingular**) if there is a matrix B of the same size, such that

$$AB = I_n \quad \text{and} \quad BA = I_n.$$

In this case we call B the **inverse** of A , and we write $A^{-1} = B$.

Theorem. If A is invertible, then $Ax = b$ has exactly one solution for every b , namely:

$$x = A^{-1}b.$$

Fact. Suppose that A and B are invertible $n \times n$ matrices.

- (1) A^{-1} is invertible and its inverse is $(A^{-1})^{-1} = A$.
- (2) AB is invertible and its inverse is $(AB)^{-1} = B^{-1}A^{-1}$.
- (3) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$.

Theorem. Let A be an $n \times n$ matrix. Here's how to compute A^{-1} .

- (1) Row reduce the augmented matrix $(A \mid I_n)$.
- (2) If the result has the form $(I_n \mid B)$, then A is invertible and $B = A^{-1}$.
- (3) Otherwise, A is not invertible.

Theorem. An $n \times n$ matrix A is invertible if and only if it is row equivalent to I_n . In this case, the sequence of row operations taking A to I_n also takes I_n to A^{-1} .

Definition. The **determinant** of a 2×2 matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is

$$\det(A) = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc.$$

Fact. If A is a 2×2 matrix, then A is invertible if and only if $\det(A) \neq 0$. In this case,

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

Definition. A transformation $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ is **invertible** if there exists another transformation $U: \mathbf{R}^n \rightarrow \mathbf{R}^n$ such that

$$T \circ U(x) = x \quad \text{and} \quad U \circ T(x) = x$$

for all x in \mathbf{R}^n . In this case we say U is the **inverse** of T , and we write $U = T^{-1}$.

Fact. A transformation T is invertible if and only if it is both one-to-one and onto.

Theorem. If T is an invertible linear transformation with matrix A , then T^{-1} is an invertible linear transformation with matrix A^{-1} .

I'll keep all of the conditions of the IMT right here, even though we don't encounter some until later:

The Invertible Matrix Theorem. Let A be a square $n \times n$ matrix, and let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the linear transformation $T(x) = Ax$. The following statements are equivalent.

- (1) A is invertible.
- (2) T is invertible.
- (3) A is row equivalent to I_n .
- (4) A has n pivots.
- (5) $Ax = 0$ has only the trivial solution.
- (6) The columns of A are linearly independent.
- (7) T is one-to-one.
- (8) $Ax = b$ is consistent for all b in \mathbf{R}^n .
- (9) The columns of A span \mathbf{R}^n .
- (10) T is onto.
- (11) A has a left inverse (there exists B such that $BA = I_n$).
- (12) A has a right inverse (there exists B such that $AB = I_n$).
- (13) A^T is invertible.
- (14) The columns of A form a basis for \mathbf{R}^n .
- (15) $\text{Col}A = \mathbf{R}^n$.
- (16) $\dim \text{Col}A = n$.

- (17) $\text{rank}A = n$.
- (18) $\text{Nul}A = \{0\}$.
- (19) $\dim \text{Nul}A = 0$.
- (20) $\det(A) \neq 0$.
- (21) *The number 0 is not an eigenvalue of A.*

CHAPTER 5

SECTIONS 5.1 AND 5.3.

Definition. The **determinant** is a function

$$\det: \{\text{square matrices}\} \longrightarrow \mathbf{R}$$

with the following **defining properties**:

- (1) $\det(I_n) = 1$
- (2) If we do a row replacement on a matrix (add a multiple of one row to another), the determinant does not change.
- (3) If we swap two rows of a matrix, the determinant scales by -1 .
- (4) If we scale a row of a matrix by k , the determinant scales by k .

Theorem. *You can use the defining properties of the determinant to compute the determinant of any matrix using row reduction.*

Magical Properties of the Determinant.

- (1) *There is one and only one function $\det: \{\text{square matrices}\} \rightarrow \mathbf{R}$ satisfying the defining properties (1)–(4).*
- (2) *A is invertible if and only if $\det(A) \neq 0$.*
- (3) *If we row reduce A without row scaling, then*

$$\det(A) = (-1)^{\#\text{swaps}} (\text{product of diagonal entries in REF})$$

- (4) *The determinant can be computed using any of the $2n$ cofactor expansions.*
- (5) $\det(AB) = \det(A)\det(B)$ and $\det(A^{-1}) = \det(A)^{-1}$
- (6) $\det(A) = \det(A^T)$
- (7) $|\det(A)|$ *is the volume of the parallelepiped defined by the columns of A.*
- (8) *If A is an $n \times n$ matrix with transformation $T(x) = Ax$, and S is a subset of \mathbf{R}^n , then the volume of $T(S)$ is $|\det(A)|$ times the volume of S. (Even for curvy shapes S.)*
- (9) *The determinant is multi-linear in the columns (or rows) of a matrix.*

SECTION 5.2.

Definition. The *ij minor* of an $n \times n$ matrix A is the $(n - 1) \times (n - 1)$ matrix A_{ij} you get by deleting the *i*th row and the *j*th column from A.

Definition. The *ij cofactor* of A is $C_{ij} = (-1)^{i+j} \det A_{ij}$.

Definition. The **determinant** of an $n \times n$ matrix A can be calculated using **cofactor expansion** along any row or column:

$$\det A = \sum_{j=1}^n a_{ij} C_{ij} \text{ for any fixed } i$$

$$\det A = \sum_{i=1}^n a_{ij} C_{ij} \text{ for any fixed } j$$

Theorem. There are special formulas for determinants of 2×2 and 3×3 matrices:

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$$

$$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33}$$

Theorem. The determinant of an upper-triangular or lower-triangular matrix is the product of the diagonal entries.

CHAPTER 6

SECTION 6.1.

Definition. Let A be an $n \times n$ matrix.

- (1) An **eigenvector** of A is a nonzero vector v in \mathbf{R}^n such that $Av = \lambda v$, for some λ in \mathbf{R} . In other words, Av is a multiple of v .
- (2) An **eigenvalue** of A is a number λ in \mathbf{R} such that the equation $Av = \lambda v$ has a nontrivial solution.

If $Av = \lambda v$ for $v \neq 0$, we say λ is the **eigenvalue for** v , and v is an **eigenvector for** λ .

Fact. The eigenvalues of a triangular matrix are the diagonal entries.

Fact. A matrix is invertible if and only if zero is not an eigenvalue.

Fact. Eigenvectors with distinct eigenvalues are linearly independent.

Definition. Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The **λ -eigenspace** of A is the set of all eigenvectors of A with eigenvalue λ , plus the zero vector:

$$\begin{aligned} \lambda\text{-eigenspace} &= \{v \text{ in } \mathbf{R}^n \mid Av = \lambda v\} \\ &= \{v \text{ in } \mathbf{R}^n \mid (A - \lambda I)v = 0\} \\ &= \text{Nul}(A - \lambda I). \end{aligned}$$

SECTION 6.2.

Definition. Let A be an $n \times n$ matrix. The **characteristic polynomial** of A is

$$f(\lambda) = \det(A - \lambda I).$$

The **characteristic equation** of A is the equation

$$f(\lambda) = \det(A - \lambda I) = 0.$$

Fact. If A is an $n \times n$ matrix, then the characteristic polynomial of A has degree n .

Fact. The roots of the characteristic polynomial (i.e., the solutions of the characteristic equation) are the eigenvalues of A .

Fact. Similar matrices have the same characteristic polynomial, hence the same eigenvalues (but different eigenvectors in general).

Definition. The **algebraic multiplicity** of an eigenvalue λ is its multiplicity as a root of the characteristic polynomial.

SECTION 6.4.

Definition. Two $n \times n$ matrices A and B are **similar** if there is an invertible $n \times n$ matrix C such that $A = CBC^{-1}$.

Definition. An $n \times n$ matrix A is **diagonalizable** if it is similar to a diagonal matrix:

$$A = PDP^{-1} \quad \text{for } D \text{ diagonal.}$$

Fact. If $A = PDP^{-1}$ for $D = \begin{pmatrix} d_{11} & 0 & \cdots & 0 \\ 0 & d_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn} \end{pmatrix}$, then

$$A^m = PD^m P^{-1} = P \begin{pmatrix} d_{11}^m & 0 & \cdots & 0 \\ 0 & d_{22}^m & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn}^m \end{pmatrix} P^{-1}.$$

The Diagonalization Theorem. An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors. In this case, $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & \cdots & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & \cdots & | \end{pmatrix} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where v_1, v_2, \dots, v_n are linearly independent eigenvectors, and $\lambda_1, \lambda_2, \dots, \lambda_n$ are the corresponding eigenvalues (in the same order).

Corollary. An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

Procedure. *How to diagonalize a matrix A :*

- (1) Find the eigenvalues of A using the characteristic polynomial.
- (2) For each eigenvalue λ of A , compute a basis \mathcal{B}_λ for the λ -eigenspace.
- (3) If there are fewer than n total vectors in the union of all of the eigenspaces \mathcal{B}_λ , then the matrix is not diagonalizable.
- (4) Otherwise, the n vectors v_1, v_2, \dots, v_n in your eigenspace bases are linearly independent, and $A = PDP^{-1}$ for

$$P = \begin{pmatrix} | & | & & | \\ v_1 & v_2 & \cdots & v_n \\ | & | & & | \end{pmatrix} \quad \text{and} \quad D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix},$$

where λ_i is the eigenvalue for v_i .

Definition. Let λ be an eigenvalue of a square matrix A . The **geometric multiplicity** of λ is the dimension of the λ -eigenspace.

Theorem. *Let λ be an eigenvalue of a square matrix A . Then*

$$1 \leq (\text{the geometric multiplicity of } \lambda) \leq (\text{the algebraic multiplicity of } \lambda).$$

Corollary. *Let λ be an eigenvalue of a square matrix A . If the algebraic multiplicity of λ is 1, then the geometric multiplicity is also 1.*

The Diagonalization Theorem (Alternate Form). *Let A be an $n \times n$ matrix. The following are equivalent:*

- (1) A is diagonalizable.
- (2) The sum of the geometric multiplicities of the eigenvalues of A equals n .
- (3) The sum of the algebraic multiplicities of the eigenvalues of A equals n , and the geometric multiplicity equals the algebraic multiplicity of each eigenvalue.

SECTION 6.5.

Review. Arithmetic in the complex numbers.

The Fundamental Theorem of Algebra. *Every polynomial of degree n has exactly n complex roots, counted with multiplicity.*

Fact. *Complex roots of real polynomials come in conjugate pairs.*

Fact. *If λ is an eigenvalue of a real matrix with eigenvector v , then $\bar{\lambda}$ is also an eigenvalue, with eigenvector \bar{v} .*

CHAPTER 7

SECTION 7.1.

Definition.

The **dot product** of two vectors x, y in \mathbf{R}^n is

$$x \cdot y = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \cdot \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} \stackrel{\text{def}}{=} x_1 y_1 + x_2 y_2 + \cdots + x_n y_n.$$

Thinking of x, y as column vectors, this is the same as the number $x^T y$.

Definition. The **length** or **norm** of a vector x in \mathbf{R}^n is

$$\|x\| = \sqrt{x \cdot x}.$$

Fact. If x is a vector and c is a scalar, then $\|cx\| = |c| \cdot \|x\|$.

Definition. The **distance** between two points x, y in \mathbf{R}^n is

$$\text{dist}(x, y) = \|y - x\|.$$

Definition. A **unit vector** is a vector v with length $\|v\| = 1$.

Definition. Let x be a nonzero vector in \mathbf{R}^n . The **unit vector in the direction of x** is the vector $x/\|x\|$.

Definition. Two vectors x, y are **orthogonal** or **perpendicular** if $x \cdot y = 0$.

Notation: $x \perp y$.

Fact. $x \perp y \iff \|x - y\|^2 = \|x\|^2 + \|y\|^2$

SECTION 7.2.

Definition. Let W be a subspace of \mathbf{R}^n . Its **orthogonal complement** is

$$W^\perp = \{v \text{ in } \mathbf{R}^n \mid v \cdot w = 0 \text{ for all } w \text{ in } W\}.$$

Fact. Let W be a subspace of \mathbf{R}^n .

- (1) W^\perp is also a subspace of \mathbf{R}^n
- (2) $(W^\perp)^\perp = W$
- (3) $\dim W + \dim W^\perp = n$
- (4) If $W = \text{Span}\{v_1, v_2, \dots, v_m\}$, then

$$\begin{aligned} W^\perp &= \text{all vectors orthogonal to each } v_1, v_2, \dots, v_m \\ &= \{x \text{ in } \mathbf{R}^n \mid x \cdot v_i = 0 \text{ for all } i = 1, 2, \dots, m\} \\ &= \text{Nul} \begin{pmatrix} -v_1^T & - \\ -v_2^T & - \\ \vdots & \\ -v_m^T & - \end{pmatrix}. \end{aligned}$$

Definition. The **row space** of an $m \times n$ matrix A is the span of the rows of A . It is denoted $\text{Row}A$. Equivalently, it is the column span of A^T :

$$\text{Row}A = \text{Col}A^T.$$

It is a subspace of \mathbf{R}^n .

Fact. $\text{Span}\{v_1, v_2, \dots, v_m\}^\perp = \text{Nul} \begin{pmatrix} -v_1^T & - \\ -v_2^T & - \\ \vdots & \\ -v_m^T & - \end{pmatrix}.$

Fact. Let A be a matrix.

- (1) $(\text{Row}A)^\perp = \text{Nul}A$ and $(\text{Nul}A)^\perp = \text{Row}A.$
- (2) $(\text{Col}A)^\perp = \text{Nul}A^T$ and $(\text{Nul}A^T)^\perp = \text{Col}A.$

SECTION 7.3.

Definition. Let $L = \text{Span}\{u\}$ be a line in \mathbf{R}^n , and let x be in \mathbf{R}^n . The **orthogonal projection of x onto L** is the point

$$\text{proj}_L(x) = \frac{x \cdot u}{u \cdot u} u.$$

Fact. Let W be a subspace of \mathbf{R}^n . Every vector x can be decomposed uniquely as

$$x = x_W + x_{W^\perp}$$

where x_W is the closest vector to x in W , and x_{W^\perp} is in W^\perp .

Theorem. Let W be a subspace of \mathbf{R}^n , and let x be a vector in \mathbf{R}^n . Then $\text{proj}_W(x)$ is the closest point to x in W . Therefore

$$x_W = \text{proj}_W(x) \quad \text{and} \quad x_{W^\perp} = x - \text{proj}_W(x).$$

Best Approximation Theorem. Let W be a subspace of \mathbf{R}^n , and let x be a vector in \mathbf{R}^n . Then $y = \text{proj}_W(x)$ is the closest point in W to x , in the sense that

$$\text{dist}(x, y') \geq \text{dist}(x, y) \quad \text{for all } y' \text{ in } W.$$

Definition. We can think of orthogonal projection as a transformation:

$$\text{proj}_W : \mathbf{R}^n \longrightarrow \mathbf{R}^n \quad x \mapsto \text{proj}_W(x).$$

Theorem. Let W be a subspace of \mathbf{R}^n .

- (1) proj_W is a linear transformation.
- (2) For every x in W , we have $\text{proj}_W(x) = x$.
- (3) For every x in W^\perp , we have $\text{proj}_W(x) = 0$.
- (4) The range of proj_W is W .

Fact. Let W be an m -dimensional subspace of \mathbf{R}^n , let $\text{proj}_W : \mathbf{R}^n \rightarrow W$ be the projection, and let A be the matrix for proj_L .

- (1) A is diagonalizable with eigenvalues 0 and 1; it is similar to the diagonal matrix with m ones and $n - m$ zeros on the diagonal.
- (2) $A^2 = A$.

SECTION 7.5.

Definition. A **least squares solution** to $Ax = b$ is a vector \hat{x} in \mathbf{R}^n such that

$$\|b - A\hat{x}\| \leq \|b - Ax\|$$

for all x in \mathbf{R}^n .

Theorem. *The least squares solutions to $Ax = b$ are the solutions to*

$$(A^T A)\hat{x} = A^T b.$$

Theorem. *If A has orthogonal columns v_1, v_2, \dots, v_n , then the least squares solution to $Ax = b$ is*

$$\hat{x} = \left(\frac{b \cdot v_1}{v_1 \cdot v_1}, \frac{b \cdot v_2}{v_2 \cdot v_2}, \dots, \frac{b \cdot v_n}{v_n \cdot v_n} \right).$$

Theorem. *Let A be an $m \times n$ matrix. The following are equivalent:*

- (1) $Ax = b$ has a unique least squares solution for all b in \mathbf{R}^m .
- (2) The columns of A are linearly independent.
- (3) $A^T A$ is invertible.

In this case, the least squares solution is $(A^T A)^{-1}(A^T b)$.

Review. Examples of best fit problems using least squares.