

## Announcements Feb 26

- Midterm 2 on **March 6**
- I will be away for the review on March 4 (Jankowski will sub)
- WeBWork 3.2 and 3.3 due Thursday
- Mid-semester evaluation under Quizzes on Canvas (due today!)
- **My office hours Monday 3-4 and Wed 2-3** in Skiles 234
- **Pop-up office hours today Wed 11-11:30 (this week only)** in Skiles 234
- TA office hours in Skiles 230 (you can go to any of these!)
  - ▶ Isabella Thu 2-3
  - ▶ Kyle Thu 1-3
  - ▶ Kalen Mon/Wed 1-1:50
  - ▶ Sidhanth Tue 10:45-11:45
- PLUS sessions Mon/Wed 6-7 LLC West with Miguel
- Supplemental problems and practice exams on the master web site

# Section 3.5

## Matrix Inverses

## Section 3.5 Outline

- The definition of a matrix inverse
- How to find a matrix inverse
- Inverses for linear transformations

## Inverses

To solve

$$Ax = b$$

we might want to “divide both sides by  $A$ ”.

We will make sense of this...

## Inverses

$A = n \times n$  matrix.

$A$  is **invertible** if there is a matrix  $B$  with

$$AB = BA = I_n$$

$B$  is called the **inverse** of  $A$  and is written  $A^{-1}$

Example:

$$\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}$$

## The $2 \times 2$ Case

Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . Then  $\det(A) = ad - bc$  is the **determinant** of  $A$ .

*Fact.* If  $\det(A) \neq 0$  then  $A$  is invertible and  $A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ .

If  $\det(A) = 0$  then  $A$  is not invertible.

*Example.*  $\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}^{-1} = -\frac{1}{2} \begin{pmatrix} 4 & -2 \\ -3 & 1 \end{pmatrix}$ .

## Solving Linear Systems via Inverses

**Fact.** If  $A$  is invertible, then  $Ax = b$  has exactly one solution:

$$x = A^{-1}b.$$

Solve

$$2x + 3y + 2z = 1$$

$$x + 3z = 1$$

$$2x + 2y + 3z = 1$$

Using

$$\begin{pmatrix} 2 & 3 & 2 \\ 1 & 0 & 3 \\ 2 & 2 & 3 \end{pmatrix}^{-1} = \begin{pmatrix} -6 & -5 & 9 \\ 3 & 2 & -4 \\ 2 & 2 & -3 \end{pmatrix}$$

## Solving Linear Systems via Inverses

What if we change  $b$ ?

$$2x + 3y + 2z = 1$$

$$x + 3z = 0$$

$$2x + 2y + 3z = 1$$

Using

$$\begin{pmatrix} 2 & 3 & 2 \\ 1 & 0 & 3 \\ 2 & 2 & 3 \end{pmatrix}^{-1} = \begin{pmatrix} -6 & -5 & 9 \\ 3 & 2 & -4 \\ 2 & 2 & -3 \end{pmatrix}$$

So finding the inverse is essentially the same as solving all  $Ax = b$  equations at once (fixed  $A$ , varying  $b$ ).



## Some Facts

Say that  $A$  and  $B$  are invertible  $n \times n$  matrices.

- $A^{-1}$  is invertible and  $(A^{-1})^{-1} = A$
- $AB$  is invertible and  $(AB)^{-1} = B^{-1}A^{-1}$

What is  $(ABC)^{-1}$ ?

## A recipe for the inverse

Suppose  $A = n \times n$  matrix.

- Row reduce  $(A | I_n)$
- If reduction has form  $(I_n | B)$  then  $A$  is invertible and  $B = A^{-1}$ .
- Otherwise,  $A$  is not invertible.

Example. Find  $\begin{pmatrix} 1 & 0 & 4 \\ 0 & 1 & 2 \\ 0 & -3 & -4 \end{pmatrix}^{-1}$

$$\begin{aligned} \left( \begin{array}{ccc|ccc} 1 & 0 & 4 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & -3 & -4 & 0 & 0 & 1 \end{array} \right) &\rightsquigarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 4 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 2 & 0 & 3 & 1 \end{array} \right) \\ &\rightsquigarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -6 & -2 \\ 0 & 1 & 0 & 0 & -2 & -1 \\ 0 & 0 & 1 & 0 & 3/2 & 1/2 \end{array} \right) \end{aligned}$$

What if you try this on one of our  $2 \times 2$  examples, such as  $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ ?

## Why Does This Work?

First answer: we can think of the algorithm as simultaneously solving

$$Ax_1 = e_1$$

$$Ax_2 = e_2$$

and so on. But the columns of  $A^{-1}$  are  $A^{-1}e_i$ , which is  $x_i$ .

## Matrix algebra with inverses

We saw that if  $Ax = b$  and  $A$  is invertible then  $x = A^{-1}b$ .

We can also, for example, solve for the matrix  $X$ , assuming that

$$AX(C + DX)^{-1} = B$$

Assume that all matrices arising in the problem are  $n \times n$  and invertible.

## Invertible Functions

A function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is **invertible** if there is a function  $U : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , so

$$T \circ U = U \circ T = \text{identity}$$

That is,

$$T \circ U(v) = U \circ T(v) = v \text{ for all } v \in \mathbb{R}^n$$

**Fact.** Suppose  $A = n \times n$  matrix and  $T$  is the matrix transformation. Then  $T$  is invertible *as a function* if and only if  $A$  is invertible. And in this case, the standard matrix for  $T^{-1}$  is  $A^{-1}$ .

**Example.** Counterclockwise rotation by  $\pi/2$ .

## Which are invertible?

Poll

Which are invertible linear transformations of  $\mathbb{R}^2$ ?

- reflection about the  $x$ -axis
- projection to the  $x$ -axis
- rotation by  $\pi$
- reflection through the origin
- a shear
- dilation by 2

## Summary of Section 3.5

- $A$  is **invertible** if there is a matrix  $B$  (called the inverse) with

$$AB = BA = I_n$$

- For a  $2 \times 2$  matrix  $A$  we have that  $A$  is invertible exactly when  $\det(A) \neq 0$  and in this case

$$A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

- If  $A$  is invertible, then  $Ax = b$  has exactly one solution:

$$x = A^{-1}b.$$

- $(A^{-1})^{-1} = A$  and  $(AB)^{-1} = B^{-1}A^{-1}$
- Recipe for finding inverse: row reduce  $(A | I_n)$ .
- Invertible linear transformations correspond to invertible matrices.