



# Announcements April 1

- Midterm 3 on **April 17**
- WeBWork 5.5 & 5.6 due Thu Apr 9.
- Quiz 8am Fri - 8am Sat on 5.2 & 5.4.  
Expect less-Googlable questions.
- My office hours Monday 3-4, Wed 2-3, and by appointment
- TA office hours on Blue Jeans (you can go to any of these!)
  - ▶ Isabella Wed 11-12
  - ▶ Kyle Wed 3-5, Thu 1-3
  - ▶ Kalen Mon/Wed 1-2
  - ▶ Sidhanth Tue 10-12
- Supplemental problems and practice exams on the master web site
- Counseling Center: <http://counseling.gatech.edu> 

# Taffy pullers

How efficient is this taffy puller?



If you run the taffy puller, the taffy starts to look like the shape on the right. Every rotation of the machine changes the number of strands of taffy by a matrix:

$$\begin{pmatrix} 1 & 0 & 2 \\ 2 & 1 & 2 \\ 4 & 2 & 3 \end{pmatrix}$$

The largest eigenvalue  $\lambda$  of this matrix describes the efficiency of the taffy puller. With every rotation, the number of strands multiplies by  $\lambda$ .

# Section 5.5

## Complex Eigenvalues

## Outline of Section 5.5

- Rotation matrices have no eigenvectors
- Crash course in complex numbers
- Finding complex eigenvectors and eigenvalues
- Complex eigenvalues correspond to rotations + dilations

▶ Demo

▶ Demo

## A matrix without an eigenvector

Recall that rotation matrices like

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$$

and

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

have no eigenvectors. Why?

# Imaginary numbers

*Problem.* When solving polynomial equations, we often run up against the issue that we can't take the square root of a negative number:

$$x^2 + 1 = 0$$

*Solution.* Take square roots of negative numbers:

$$x = \pm\sqrt{-1}$$

We usually write  $\sqrt{-1}$  as  $i$  (for “imaginary”), so  $x = \pm i$ .

Now try solving these:

$$x^2 + 3 = 0$$

$$x^2 - x + 1 = 0$$

# Complex numbers

We can add/multiply (and divide!) complex numbers:

$$(2 - 3i) + (-1 + i) =$$

$$(2 - 3i)(-1 + i) =$$

# Complex numbers

The complex numbers are the numbers

$$\mathbb{C} = \{a + bi \mid a, b \text{ in } \mathbb{R}\}$$

We can **conjugate** complex numbers:  $\overline{a + bi} = a - bi$



# Complex numbers and polynomials

**Fundamental theorem of algebra.** Every polynomial of degree  $n$  has exactly  $n$  complex roots (counted with multiplicity).

**Fact.** If  $z$  is a root of a real polynomial then  $\bar{z}$  is also a root.

So what are the possibilities for degree 2, 3 polynomials?

# Complex eigenvalues

Say  $A$  is a square matrix with real entries.

We can now find **complex** eigenvectors and eigenvalues.

**Fact.** If  $\lambda$  is an eigenvalue of  $A$  with eigenvector  $v$  then  $\bar{\lambda}$  is an eigenvalue of  $A$  with eigenvector  $\bar{v}$ .

Why?

## Complex eigenvalues

Find the complex eigenvalues and eigenvectors for

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

## Three shortcuts for complex eigenvectors

Suppose we have a  $2 \times 2$  matrix with complex eigenvalue  $\lambda$ .

(1) We do not need to row reduce  $A - \lambda I$  by hand; we know the bottom row will become zero.

(2) Then if the reduced matrix is:

$$A = \begin{pmatrix} x & y \\ 0 & 0 \end{pmatrix}$$

the eigenvector is

$$A = \begin{pmatrix} -y \\ x \end{pmatrix}$$

(3) Also, we get the other eigenvalue/eigenvector pair for free: conjugation.

## Complex eigenvalues

Find the complex eigenvalues and eigenvectors for

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & -2 \\ 1 & 3 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -2 \\ 0 & 2 & 0 \end{pmatrix}$$

## Review for Section 5.5

If  $v$  is an eigenvector for  $A$  with complex entries, is  $i \cdot v$  also an eigenvector for  $A$ ?

If  $A$  is a  $4 \times 4$  matrix with real entries, what are the possibilities for the number of non-real eigenvalues of  $A$ ?

## Summary of Section 5.5

- Complex numbers allow us to solve all polynomials completely, and find  $n$  eigenvectors for an  $n \times n$  matrix
- If  $\lambda$  is an eigenvalue with eigenvector  $v$  then  $\bar{\lambda}$  is an eigenvalue with eigenvector  $\bar{v}$