

# ABSTRACT COMMENSURATORS OF RIGHT-ANGLED ARTIN GROUPS AND MAPPING CLASS GROUPS

MATT CLAY, CHRISTOPHER J. LEININGER, AND DAN MARGALIT

ABSTRACT. We prove that, aside from the obvious exceptions, the mapping class group of a compact orientable surface is not abstractly commensurable with any right-angled Artin group. Our argument applies to various subgroups of the mapping class group—the subgroups generated by powers of Dehn twists and the terms of the Johnson filtration—and additionally to the outer automorphism group of a free group and to certain linear groups.

There are many analogies and interconnections between the theories of right-angled Artin groups on one hand and mapping class groups on the other hand. For instance, by the work of Crisp and Wiest [12], the work of Koberda [26], and the work of the first two authors with Mangahas [11], there is an abundance of injective homomorphisms from right-angled Artin groups to mapping class groups. Also, the last two authors proved [29] that any two elements of the pure braid group either generate a free group or a free abelian group—a property shared by all right-angled Artin groups. We are thus led to ask to what extent mapping class groups are the same as right-angled Artin groups.

It is straightforward to see that most mapping class groups are not isomorphic to right-angled Artin groups, for instance because right-angled Artin groups are torsion free. On the other hand, mapping class groups have finite-index subgroups that are torsion free, and so this leaves open the possibility that mapping class groups are abstractly commensurable to right-angled Artin groups, that is, that they have isomorphic finite-index subgroups. We prove that, aside from a small number of exceptions, this is not the case. We also extend this result to several classes of groups related to mapping class groups. We start by recalling some definitions.

To a finite graph  $\Gamma$ , we can associate a right-angled Artin group: this is the group with one generator for each vertex of  $\Gamma$ , and one defining relator for each edge, namely, the commutator of the two generators corresponding to the endpoints.

Let  $S_{g,n}$  denote a closed, connected, orientable surface of genus  $g$  with  $n$  marked points. The mapping class group  $\text{Mod}(S_{g,n})$  is the group of homotopy classes of orientation-preserving homeomorphisms of  $S_{g,n}$  preserving the marked points.

As discussed in Koberda’s paper [26, Theorem 1.5], no finite-index subgroup of  $\text{Mod}(S_{g,n})$  injects into a right-angled Artin group if  $g \geq 2$  and  $(g, n) \neq (2, 0)$ ; see also [23]. In particular, such mapping class groups are not abstractly commensurable

---

2000 *Mathematics Subject Classification*. Primary: 20E36; Secondary: 57M07.

*Key words and phrases*. right-angled Artin group, mapping class group, abstract commensurator.

The authors gratefully acknowledge support from the National Science Foundation.

with right-angled Artin groups. The last statement has a quick proof: for a right-angled Artin group, the virtual cohomological dimension is equal to the maximal rank of a free abelian subgroup, while for mapping class groups these numbers—which are invariant under passage to finite-index subgroups—are equal if and only if  $g = 0$ ,  $g = 1$ , or  $(g, n) = (2, 0)$ ; see [22, Theorem 6.4.A–C] and [7, Theorem A].

As for the other mapping class groups, the first two authors proved with Mangahas that  $\text{Mod}(S_{2,0})$  is not abstractly commensurable with any right-angled Artin group [11, Proposition 7.2] and explained how to apply their method to the case of  $\text{Mod}(S_{0,n})$  with  $n \geq 6$ . Our first theorem extends these results to the remaining groups  $\text{Mod}(S_{g,n})$ , while at the same time giving a uniform argument for all cases.

**Theorem 1.** *Let  $g, n \geq 0$  and assume that  $3g + n \geq 5$ . No right-angled Artin group is abstractly commensurable with  $\text{Mod}(S_{g,n})$ .*

To prove Theorem 1, we consider a third invariant of the abstract commensurability class of a group  $G$ , namely, the abstract commensurator  $\text{Comm}(G)$ . This is the group of all isomorphisms between finite-index subgroups of  $G$ , up to restriction; see [21, Section 5]. For most of the surfaces covered by Theorem 1, Ivanov [21, Theorem 5] and Korkmaz [27, Theorem 3] showed that  $\text{Comm}(\text{Mod}(S_{g,n}))$  is isomorphic to the extended mapping class group  $\text{Mod}^\pm(S_{g,n})$ , the group of homotopy classes of all (not-necessarily-orientation-preserving) homeomorphisms of  $S_{g,n}$ . There are two exceptions:  $\text{Comm}(\text{Mod}(S_{1,2})) \cong \text{Mod}^\pm(S_{0,5})$  and  $\text{Comm}(\text{Mod}(S_{2,0})) \cong \text{Mod}^\pm(S_{0,6})$ ; see [5, Proposition 7] and [20, Theorem 1.2].

Let  $A$  be an arbitrary right-angled Artin group; we must show that  $A$  is not abstractly commensurable with  $\text{Mod}(S_{g,n})$  when  $3g + n \geq 5$ . We may assume  $A$  is not isomorphic to  $\mathbb{Z}^m$ . Indeed, in this case  $A$  is not abstractly commensurable with  $\text{Mod}(S_{g,n})$ , for instance because nonzero powers of Dehn twists about nondisjoint curves fail to commute [18, Section 3.3]. Applying the previous paragraph, it remains to show that if  $A$  is not free abelian, then  $\text{Comm}(A)$  is not isomorphic to any  $\text{Mod}^\pm(S_{g,n})$  with  $3g + n \geq 5$ . To do this, we will show that  $\text{Comm}(A)$  contains  $(\mathbb{Z}/2\mathbb{Z})^k$  for arbitrarily large  $k$ ; it follows from Kerckhoff’s solution to the Nielsen realization problem [24] that  $\text{Mod}^\pm(S_{g,n})$  does not contain finite subgroups of arbitrary large cardinality.

The proof of the fact that  $(\mathbb{Z}/2\mathbb{Z})^k \leq \text{Comm}(A)$  for large  $k$  consists of two observations. The first observation is that if the defining graph  $\Gamma$  for  $A$  has  $k$  vertices, then  $\text{Comm}(A)$  contains a subgroup isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^k$ . The generators are the abstract commensurators obtained by inverting the elements of  $A$  corresponding to the vertices of  $\Gamma$  (each is a nontrivial element of  $\text{Comm}(A)$  since every finite-index subgroup of  $A$  contains some power of this generator). The second observation is that (as  $A$  is not free abelian)  $A$  contains a finite-index subgroup  $A'$  that is a right-angled Artin group whose defining graph has arbitrarily many vertices. Since  $\text{Comm}(A) \cong \text{Comm}(A')$ , the result follows.

To prove the second observation, we choose a vertex  $v$  of  $\Gamma$  for which the star of  $v$  is not all of  $\Gamma$ . Such a vertex exists since  $A$  is not free abelian. Consider the homomorphism  $A \rightarrow \mathbb{Z}/m\mathbb{Z}$  obtained by sending the generator of  $A$  corresponding

to  $v$  to 1 and the generators corresponding to all other vertices to 0. The kernel has finite index in  $A$  and is a right-angled Artin group whose defining graph is obtained by taking  $m$  copies of  $\Gamma$  and gluing along the  $m$  copies of the star of  $v$ ; see [6, Section 11] and [25, Corollary 5]. Since the star of  $v$  is not all of  $\Gamma$ , the new graph has more than  $m$  vertices. This completes the proof.

Our construction of  $(\mathbb{Z}/2\mathbb{Z})^k$  subgroups of  $\text{Comm}(A)$  can be combined with a theorem of Bartholdi and Bogopolski [3, Theorem 2.8] to prove that  $\text{Comm}(A)$  is not finitely generated, thus giving a different (but similar) proof that  $\text{Comm}(A)$  is not isomorphic to any  $\text{Mod}^\pm(S)$ .

The assumptions in Theorem 1 are in fact necessary, as  $\text{Mod}(S_{1,0}) \cong \text{Mod}(S_{1,1}) \cong \text{SL}_2(\mathbb{Z})$  and  $\text{Mod}(S_{0,4}) \cong \text{PSL}_2(\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$  are commensurable with the free group  $F_2$  and  $\text{Mod}(S_{0,n})$  is finite, hence abstractly commensurable with the trivial right-angled Artin group, for  $n \leq 3$ .

Finally, the  $(\mathbb{Z}/2\mathbb{Z})^k$  subgroups of  $\text{Comm}(A)$  we construct further embed into the quasi-isometry group of  $A$ , and so using the theorem of Behrstock, Kleiner, Minsky, and Mosher that the quasi-isometry group of  $\text{Mod}(S_{g,n})$  is again an extended mapping class group [4, Theorem 1.1], we can conclude the strengthening of Theorem 1 that most mapping class groups are not quasi-isometric to right-angled Artin groups.

**Mapping class groups of surfaces with boundary.** Let  $S_{g,n}^b$  denote the surface obtained from  $S_{g,n}$  by removing the interiors of  $b$  disks, disjoint from each other and the marked points; we denote  $S_{g,0}^b$  by  $S_g^b$  and  $S_{g,0}^0$  by  $S_g$ . The mapping class group  $\text{Mod}(S_{g,n}^b)$  is the group of homotopy classes of orientation-preserving homeomorphisms of  $S_{g,n}^b$  that restrict to the identity on the boundary.

**Theorem 2.** *Let  $g, n, b \geq 0$  and assume that  $3g + n + b \geq 5$ . No right-angled Artin group is abstractly commensurable with  $\text{Mod}(S_{g,n}^b)$ .*

As the braid group on  $n$  strands is isomorphic to  $\text{Mod}(S_{0,n}^1)$ , Theorem 2 in particular implies that the braid group (or pure braid group) on  $n \geq 4$  strands is not abstractly commensurable with any right-angled Artin group.

When  $b > 0$ , the virtual cohomological dimension of  $\text{Mod}(S_{g,n}^b)$  and the maximal rank of an abelian subgroup of  $\text{Mod}(S_{g,n}^b)$  are equal if and only if  $g \in \{0, 1\}$ , and so it suffices to consider these two values of  $g$ . We will use a property special to these two cases, namely, that  $\text{Mod}(S_{g,n}^b)$  is abstractly commensurable with the direct product  $\text{Mod}(S_{g,n+b}) \times \mathbb{Z}^b$ ; see Theorem 8 below.

If a right-angled Artin group  $A$  is abstractly commensurable with  $\text{Mod}(S_{g,n}^b)$ , it is then abstractly commensurable with  $\text{Mod}(S_{g,n+b}) \times \mathbb{Z}^b$ . From this it follows that  $A$  is isomorphic to  $A' \times \mathbb{Z}^b$  for some right-angled Artin group  $A'$  (use [31, Section III]) and  $A'$  is abstractly commensurable with  $\text{Mod}(S_{g,n+b})$  (cf. [28, Lemma 2.2]), violating Theorem 1.

In principle, Theorem 2 can be proven in the same way as Theorem 1. However when  $b > 0$  the group  $\text{Comm}(\text{Mod}(S_{g,n}^b))$  has only been computed for  $g = 0$  and  $b =$

1; see [28]. Using Theorem 8 below, Section 3 of [28], and the fact that  $\text{Mod}(S_{1,n+b})$  surjects onto the virtually free group  $\text{Mod}(S_{1,1}) \cong \text{SL}_2(\mathbb{Z})$ , though, it follows that  $\text{Comm}(\text{Mod}(S_{g,n}^b)) \cong \text{Comm}(\text{Mod}(S_{g,n+b})) \times (\text{GL}_b(\mathbb{Q}) \times (\mathbb{Q}^b)^\infty)$  for  $g = 0, 1$ .

**Subgroups of the mapping class group.** Our proof of Theorem 1 shows that if  $G$  is any group that is not virtually abelian and has the property that  $\text{Comm}(G)$  does not contain finite subgroups of arbitrarily large cardinality, then  $G$  is not abstractly commensurable with any right-angled Artin group. We will apply this criterion in several other cases; first for certain subgroups of the mapping class group, and then for the outer automorphism group of a free group and certain linear groups.

Let  $\mathcal{T}(S_g, d)$  denote the subgroup of  $\text{Mod}(S_g)$  generated by the  $d$ th powers of all Dehn twists. This group has infinite index in  $\text{Mod}(S_g)$  for  $d \geq 11$  and  $g \geq 2$  [19]. It has been conjectured that  $\mathcal{T}(S_g, d)$  is a right-angled Artin group for  $d$  large [19]. However, Ivanov's proof that  $\text{Comm}(\text{Mod}(S_g)) \cong \text{Mod}^\pm(S_g)$  carries over directly to show that  $\text{Comm}(\mathcal{T}(S_g, d)) \cong \text{Mod}^\pm(S_g)$  for  $g \geq 3$ ; see [1] for details. As  $\mathcal{T}(S_g, d)$  is not virtually abelian, we conclude the following.

**Theorem 3.** *Let  $g \geq 3$  and  $d \geq 1$ . The group  $\mathcal{T}(S_g, d)$  is not abstractly commensurable with any right-angled Artin group.*

Let  $\pi$  denote  $\pi_1(S_g)$ , and let  $\pi^k$  denote the  $k$ th term of its lower central series:  $\pi^1 = \pi$  and  $\pi^{k+1} = [\pi, \pi^k]$ . The Johnson filtration of  $\text{Mod}(S_g)$  is the nested sequence of groups  $(\mathcal{N}_k(S_g))$  where  $\mathcal{N}_k(S_g)$  is the kernel of the natural homomorphism  $\text{Mod}(S_g) \rightarrow \text{Out}(\pi/\pi^{k+1})$ . The intersection of the  $\mathcal{N}_k(S_g)$  is trivial. The groups  $\mathcal{N}_1(S_g)$  and  $\mathcal{N}_2(S_g)$  are also known as the Torelli group and Johnson kernel of  $S_g$ . For  $g$  large enough, the abstract commensurators of these groups are all known to be isomorphic to  $\text{Mod}^\pm(S_g)$  [17, Theorem 7] [9, Main Theorem 1] [10].

**Theorem 4.** *Let  $k \geq 1$ . Let  $g \geq 3$  if  $k \leq 3$  and let  $g \geq 4$  if  $k > 3$ . No right-angled Artin group is abstractly commensurable with  $\mathcal{N}_k(S_g)$ .*

**Outer automorphism groups of right-angled Artin groups.** For  $g \geq 1$ , the group  $\text{Mod}^\pm(S_g)$  is isomorphic to  $\text{Out}(\pi_1(S_g))$ . As  $\pi_1(S_1) \cong \mathbb{Z}^2$  and as  $\pi_1(S_{g,n})$  is a free group for  $n > 0$ , we can obtain analogs of the mapping class group by considering groups of the form  $\text{Out}(A)$  where  $A$  is a right-angled Artin group. By results of Davis–Januszkiewicz [14] and Taylor [33], there are again many embeddings of right-angled Artin groups into  $\text{Out}(A)$  where  $A$  is  $\mathbb{Z}^n$  or  $F_n$ .

By the combined work of Borel [8] and Margulis [30], the abstract commensurator of  $\text{GL}_n(\mathbb{Z}) \cong \text{Out}(\mathbb{Z}^n)$  is  $\text{PGL}_n(\mathbb{Q}) \times \mathbb{Z}/2\mathbb{Z}$  for  $n \geq 3$ ; see e.g. [32, Section 7.3] for an exposition of these ideas. Since  $\text{PGL}_n(\mathbb{Q})$  does not contain arbitrarily large subgroups of the form  $(\mathbb{Z}/2\mathbb{Z})^k$  and since  $\text{GL}_n(\mathbb{Z})$  is not virtually abelian, we obtain the following.

**Theorem 5.** *Let  $n \geq 3$ . No right-angled Artin group is abstractly commensurable with  $\text{Out}(\mathbb{Z}^n) \cong \text{GL}_n(\mathbb{Z})$ .*

Wortman [34] has pointed out that the argument for Theorem 5 extends to any lattice in a semisimple Lie group not locally isomorphic to  $\text{SL}_2(\mathbb{R})$ . Indeed, for any

such lattice  $G$ , Mostow–Prasad–Margulis superrigidity implies that  $\text{Comm}(G)$  is a subgroup of some  $\text{GL}_n(\mathbb{C})$ . From Jordan canonical form, we know that  $\text{GL}_n(\mathbb{C})$  contains subgroups isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^k$  only for  $k \leq n$ . In particular, we can deduce the theorem of Koberda that for  $k \geq 3$  no right-angled Artin group is abstractly commensurable with a lattice in  $\text{SO}(k, 1)$  [26, Theorem 1.14]. Additionally, Studenmund [32] has shown that the abstract commensurators of many lattices with nontrivial solvable radicals (e.g.  $\text{SL}_n(\mathbb{Z}) \rtimes \mathbb{Z}^n$ ) are linear, so our argument applies to these lattices as well.

Farb and Handel proved that  $\text{Comm}(\text{Out}(F_n))$  is isomorphic to  $\text{Out}(F_n)$  when  $n \geq 4$  (it is not known whether  $\text{Comm}(\text{Out}(F_3)) \cong \text{Out}(F_3)$  or not) [16]. A finite subgroup of  $\text{Out}(F_n)$  can be identified with the symmetries of some fixed metric graph of rank  $n$  [13, Theorem 2.1] [35], and so there is a bound on the cardinality of such subgroups that depends only on  $n$ . Since  $\text{Out}(F_n)$  is not virtually abelian, we obtain the following.

**Theorem 6.** *Let  $n \geq 4$ . No right-angled Artin group is abstractly commensurable with  $\text{Out}(F_n)$ .*

Theorems 5 and 6 tell us that (in most cases) the outer automorphism groups of the right-angled Artin groups  $\mathbb{Z}^n$  and  $F_n$  are not commensurable with right-angled Artin groups. We also know that  $\text{Out}(\mathbb{Z}^2) \cong \text{Out}(F_2) \cong \text{GL}_2(\mathbb{Z})$  is commensurable with  $F_2$ , and that the outer automorphism group of the right-angled Artin group associated to the linear graph with four vertices is  $((\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})^4) \times \mathbb{Z}^4$  (cf. [15, Proposition 2.15]), which is abstractly commensurable with  $\mathbb{Z}^4$ . We are therefore led to the following question.

**Question 7.** *For which right-angled Artin groups  $A$  is  $\text{Out}(A)$  abstractly commensurable with a right-angled Artin group? Can we have  $\text{Out}(A)$  abstractly commensurable with a right-angled Artin group that is not free or free abelian?*

APPENDIX. VIRTUAL SPLITTING FOR LOW-GENUS MAPPING CLASS GROUPS

In this appendix we prove the following fact, used in our proof of Theorem 2. For the statement, we define the level two subgroup  $\text{Mod}(S_{g,n}^b)[2]$  of  $\text{Mod}(S_{g,n}^b)$  as the finite-index subgroup consisting of all elements that act trivially on the mod two homology of the surface obtained from  $S_{g,n}^b$  by removing the marked points. Note that when  $g = 0$  the group  $\text{Mod}(S_{0,n}^b)[2]$  is the same the pure mapping class group; in particular  $\text{Mod}(S_{0,n}^1)[2]$  is isomorphic to the pure braid group on  $n$  strands.

**Theorem 8.** *Let  $g \in \{0, 1\}$ , and assume  $3g + n + b \geq 3$ . We have*

$$\text{Mod}(S_{g,n}^b)[2] \cong \text{Mod}(S_{g,n+b})[2] \times \mathbb{Z}^b.$$

It suffices to show that  $\text{Mod}(S_{g,n}^b)[2]$  splits as a direct product over its center, the free abelian group generated by the Dehn twists about the  $b$  boundary components of  $S_{g,n}^b$ . The theorem then follows from the fact that the natural inclusion  $S_{g,n}^b \rightarrow S_{g,n+b}$  induces a well-defined surjective map of level two mapping class groups, and that the kernel of this map is the center [18, Proposition 3.19].

First we deal with the case  $g = 1$ . The group  $\text{Mod}(S_1^1)[2]$  is isomorphic to the pure braid group on three strands (combine [18, Section 9.4.1] with [2]). It is well-known that the latter splits over its infinite cyclic center [18, Section 9.3], and so we are done in this case. Denote by  $s$  a retraction  $\text{Mod}(S_1^1)[2] \rightarrow Z(\text{Mod}(S_1^1)[2])$  that defines the splitting.

Let  $f_i : \text{Mod}(S_{g,n}^b)[2] \rightarrow \text{Mod}(S_1^1)[2]$  be the homomorphism obtained by forgetting the  $n$  marked points and by capping each boundary component of  $S_{g,n}^b$ —except for the  $i$ th—with a disk. Also, let  $h_i : Z(\text{Mod}(S_1^1)[2]) \rightarrow \text{Mod}(S_{g,n}^b)[2]$  be the homomorphism that maps the Dehn twist about the boundary of  $S_1^1$  to the Dehn twist about the  $i$ th boundary component of  $S_{g,n}^b$ . The product of the maps  $h_i \circ s \circ f_i$  is the desired retraction  $\text{Mod}(S_{g,n}^b)[2] \rightarrow Z(\text{Mod}(S_{g,n}^b)[2])$ .

The genus zero version is nearly identical. The role of  $\text{Mod}(S_1^1)[2]$  is played by  $\text{Mod}(S_{0,2}^1)[2] \cong \mathbb{Z}$ . The only essential difference in this case is that there are many choices of homomorphism  $\text{Mod}(S_{0,n}^b)[2] \rightarrow \text{Mod}(S_{0,2}^1)[2]$  available for each coordinate of the splitting.

In contrast to Theorem 8, it is known that  $\text{Mod}(S_g^1)$  does not have a finite-index subgroup that splits over its center when  $g \geq 2$ ; see [18, Proposition 5.10].

**Acknowledgments.** We would like to thank Jason Behrstock, Matt Day, Benson Farb, Thomas Koberda, Daniel Studenmund, and Kevin Wortman for helpful conversations.

## REFERENCES

- [1] Javier Aramayona and Juan Souto. A remark on homomorphisms from right-angled artin groups to mapping class groups, 2013.
- [2] V. I. Arnol'd. A remark on the branching of hyperelliptic integrals as functions of the parameters. *Funkcional. Anal. i Priložen.*, 2(3):1–3, 1968.
- [3] L. Bartholdi and O. Bogopolski. On abstract commensurators of groups. *J. Group Theory*, 13(6):903–922, 2010.
- [4] Jason Behrstock, Bruce Kleiner, Yair Minsky, and Lee Mosher. Geometry and rigidity of mapping class groups. *Geom. Topol.*, 16(2):781–888, 2012.
- [5] Jason Behrstock and Dan Margalit. Curve complexes and finite index subgroups of mapping class groups. *Geom. Dedicata*, 118:71–85, 2006.
- [6] Mladen Bestvina, Bruce Kleiner, and Michah Sageev. The asymptotic geometry of right-angled Artin groups. I. *Geom. Topol.*, 12(3):1653–1699, 2008.
- [7] Joan S. Birman, Alex Lubotzky, and John McCarthy. Abelian and solvable subgroups of the mapping class groups. *Duke Math. J.*, 50(4):1107–1120, 1983.
- [8] Armand Borel. Density and maximality of arithmetic subgroups. *J. Reine Angew. Math.*, 224:78–89, 1966.
- [9] Tara E. Brendle and Dan Margalit. Commensurations of the Johnson kernel. *Geom. Topol.*, 8:1361–1384 (electronic), 2004.
- [10] Martin Bridson, Alexandra Pettet, and Juan Souto. Work in progress.
- [11] Matt T. Clay, Christopher J. Leininger, and Johanna Mangahas. The geometry of right-angled Artin subgroups of mapping class groups. *Groups Geom. Dyn.*, 6(2):249–278, 2012.
- [12] John Crisp and Bert Wiest. Quasi-isometrically embedded subgroups of braid and diffeomorphism groups. *Trans. Amer. Math. Soc.*, 359(11):5485–5503, 2007.

- [13] Marc Culler. Finite groups of outer automorphisms of a free group. In *Contributions to group theory*, volume 33 of *Contemp. Math.*, pages 197–207. Amer. Math. Soc., Providence, RI, 1984.
- [14] Michael W. Davis and Tadeusz Januszkiewicz. Right-angled Artin groups are commensurable with right-angled Coxeter groups. *J. Pure Appl. Algebra*, 153(3):229–235, 2000.
- [15] Matthew B. Day. On solvable subgroups of automorphism groups of right-angled Artin groups. *Internat. J. Algebra Comput.*, 21(1-2):61–70, 2011.
- [16] Benson Farb and Michael Handel. Commensurations of  $\text{Out}(F_n)$ . *Publ. Math. Inst. Hautes Études Sci.*, (105):1–48, 2007.
- [17] Benson Farb and Nikolai V. Ivanov. The Torelli geometry and its applications: research announcement. *Math. Res. Lett.*, 12(2-3):293–301, 2005.
- [18] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [19] Louis Funar. On power subgroups of mapping class groups, 2009.
- [20] Elmas Irmak. Complexes of nonseparating curves and mapping class groups. *Michigan Math. J.*, 54(1):81–110, 2006.
- [21] Nikolai V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *Internat. Math. Res. Notices*, (14):651–666, 1997.
- [22] Nikolai V. Ivanov. Mapping class groups. In *Handbook of geometric topology*, pages 523–633. North-Holland, Amsterdam, 2002.
- [23] Michael Kapovich and Bernhard Leeb. Actions of discrete groups on nonpositively curved spaces. *Math. Ann.*, 306(2):341–352, 1996.
- [24] Steven P. Kerckhoff. The Nielsen realization problem. *Ann. of Math. (2)*, 117(2):235–265, 1983.
- [25] Sang-hyun Kim and Thomas Koberda. Embedability between right-angled Artin groups. *Geom. Topol.*, 17(1):493–530, 2013.
- [26] Thomas Koberda. Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups. *Geom. Funct. Anal.*, 22(6):1541–1590, 2012.
- [27] Mustafa Korkmaz. Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology Appl.*, 95(2):85–111, 1999.
- [28] Christopher J. Leininger and Dan Margalit. Abstract commensurators of braid groups. *J. Algebra*, 299(2):447–455, 2006.
- [29] Christopher J. Leininger and Dan Margalit. Two-generator subgroups of the pure braid group. *Geom. Dedicata*, 147:107–113, 2010.
- [30] G. A. Margulis. Non-uniform lattices in semisimple algebraic groups. In *Lie groups and their representations (Proc. Summer School on Group Representations of the Bolyai János Math. Soc., Budapest, 1971)*, pages 371–553. Halsted, New York, 1975.
- [31] Herman Servatius. Automorphisms of graph groups. *J. Algebra*, 126(1):34–60, 1989.
- [32] Daniel Studenmund. Abstract commensurators of lattices in lie groups, 2013.
- [33] Samuel J. Taylor. Right-angled Artin groups and  $\text{Out}(F_n)$  I: quasi-isometric embeddings, 2013.
- [34] Kevin Wortman. Personal communication, 2013.
- [35] Bruno Zimmermann. Über Homöomorphismen  $n$ -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen. *Comment. Math. Helv.*, 56(3):474–486, 1981.

MATT CLAY, DEPARTMENT OF MATHEMATICS, 301 SCEN - 1 UNIVERSITY OF ARKANSAS, FAYETTEVILLE, AR 72701, MATTCCLAY@UARK.EDU

CHRISTOPHER J. LEININGER, DEPT. OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, 273 ALTGELD HALL, 1409 W. GREEN ST., URBANA, IL 61802, CLEIN@MATH.UIUC.EDU

DAN MARGALIT, SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, 686 CHERRY ST., ATLANTA, GA 30332, MARGALIT@MATH.GATECH.EDU